

Reanalyzing the visible colors of Centaurs and KBOs: what is there and what we might be missing.

Nuno Peixinho¹, Audrey Delsanti², and Alain Doressoundiram³

¹ Unidad de Astronomía, Fac. de Ciencias Básicas, Universidad de Antofagasta, Avda. U. de Antofagasta 02800, Antofagasta, Chile
e-mail: nuno.peixinho@uantof.cl

² Aix Marseille Université, CNRS, LAM (Laboratoire d'Astrophysique de Marseille) UMR 7326, 13388 Marseille, France
e-mail: Audrey.Delsanti@lam.fr

³ Observatoire de Paris, Site de Meudon, 5 place Jules Janssen, 92190 Meudon, France
e-mail: Alain.Doressoundiram@obspm.fr

Received 30 November 2014 / Accepted 6 February 2015

ABSTRACT

Since the discovery of the Kuiper Belt, broadband surface colors were thoroughly studied as a first approximation to the object reflectivity spectra. Visible colors (BVRI) have proven to be a reasonable proxy for real spectra, which are rather linear in this range. In contrast, near-IR colors (JHK bands) could be misleading when absorption features of ices are present in the spectra. Although the physical and chemical information provided by colors are rather limited, broadband photometry remains the best tool for establishing the bulk surface properties of KBOs and Centaurs. In this work, we explore for the first time general, recurrent effects in the study of visible colors that could affect the interpretation of the scientific results: i) how a correlation could be missed or weakened as a result of the data error bars, ii) the “risk” of missing an existing trend because of low sampling, and the possibility of making quantified predictions on the sample size needed to detect a trend at a given significance level — assuming the sample is unbiased, iii) the use of partial correlations to distinguish the mutual effect of two or more (physical) parameters, and iv) the sensitivity of the “reddening line” tool to the central wavelength of the filters used. To illustrate and apply these new tools, we have compiled the visible colors and orbital parameters of about 370 objects available in the literature — assumed, by default, as unbiased samples — and carried out a traditional analysis per dynamical family. Our results show in particular how a) data error bars impose a limit on the detectable correlations regardless of sample size and that therefore, once that limit is achieved, it is important to diminish the error bars, but it is pointless to enlarge the sampling with the same or larger errors; b) almost all dynamical families still require larger samplings to ensure the detection of correlations stronger than ± 0.5 , that is, correlations that may explain $\sim 25\%$ or more of the color variability; c) the correlation strength between (V-R) vs. (R-I) is systematically lower than the one between (B-V) vs. (V-R) and is not related with error-bar differences between these colors; d) it is statistically equivalent to use any of the different flavors of orbital excitation or collisional velocity parameters regarding the famous color-inclination correlation among classical KBOs — which no longer appears to be a strong correlation — whereas the inclination and Tisserand parameter relative to Neptune cannot be separated from one another; and e) classical KBOs are the only dynamical family that shows neither (B-V) vs. (V-R) nor (V-R) vs. (R-I) correlations. It therefore is the family with the most unpredictable visible surface reflectivities.

Key words. Kuiper belt: general — Methods: data analysis — Methods: statistical — Techniques: photometric

1. Introduction

Kuiper Belt objects (KBOs), also known as Trans-Neptunian objects (TNOs), are a large population of icy bodies orbiting the Sun with semi-major axes greater than that of Neptune. They are considered as the least altered objects in the solar system. Their existence had been hypothesized by Leonard (1930), Edgeworth (1943, 1949), and Kuiper (1951) — and the expression Edgeworth-Kuiper Belt is also used in some works — but they were only observed in 1992 by Jewitt & Luu (1993). Currently, there are more than 1700 objects identified, $\sim 4.5\%$ are binary or multiple systems, and there are probably $\sim 200\,000$ objects larger than $100\,km$ in diameter (Petit et al. 2011).

KBOs subdivide into several dynamical families, although there is no strict definition regarding the boundaries between most of them (see Gladman et al. 2008, for a review). Centaurs are, presumably, former KBOs that evolved into inner orbits, that is, both their semi-major axes and perihelia lie between the orbits of Jupiter and Neptune (Fernandez 1980; Lev-

ison & Duncan 1997) — although when using a less strict dynamical definition of Centaurs, some may have an Oort Cloud origin (Emel’yanenko et al. 2013; Fouchard et al. 2014). Every $\sim 125\,yr$ a new Centaur escapes from the Kuiper Belt into a short-lived chaotic orbit, with an average half-life of $\sim 2.75\,Myr$ — although some may last for millions of years — before becoming a Jupiter-family comet (JFC) or being re-injected into the outer solar system (e.g., Horner et al. 2004). Given this close link, Centaurs have usually been studied as one more family of KBOs.

Since the first analysis by Luu & Jewitt (1996) of a sample of three Centaurs and nine KBOs, any relation between the surface properties observed at different wavelengths has been searched for, such as different colors. These relations might be among the surface properties themselves or between the surface properties and the orbital parameters. Although correlation analysis does not allow us to predict specific values of unmeasured properties by knowing others, nor to draw conclusions about cause and

effect relationships, it is the most elementary tool for detecting and quantifying which variables or properties covary with each other, which provides a first insight into the phenomena we are observing.

Many works reported trends, patterns, and correlations between different surface colors and between those and the orbital properties of these objects (see review Doressoundiram et al. 2008). While several teams had their own surveys — for example, the Tegler, Romanishin, and Consolmagno survey (e.g., Tegler & Romanishin 1998, 2000, 2003; Tegler et al. 2003), the ESO Large Program on Centaurs and TNOs (e.g., Delsanti et al. 2001; Boehnhardt et al. 2002; Delsanti et al. 2004; Peixinho et al. 2004; Delsanti et al. 2006), the Meudon Multicolor Survey (2MS), (e.g., Barucci et al. 1999, 2000; Doressoundiram et al. 2001, 2002, 2005b, 2007), and the Second ESO Large Program (e.g., DeMeo et al. 2009; Perna et al. 2010, and references therein) — focusing on the analysis of their own datasets for methodological reasons, Hainaut & Delsanti (2002) made the first thorough analysis of a large compilation of all published data, known as the MBOSS database, which has recently been updated (Hainaut et al. 2012).

Although the physical and chemical information provided by the visible colors of these objects is rather limited, the analysis of visible colors alone has led to very rich debates on the Kuiper Belt science.

Given the narrow temperature differences in the Kuiper Belt region, KBOs were initially presumed to have a primordial homogeneous composition, and the first modeling works attempted to explain the visible surface color diversity as a result of a competition between a reddening space-weather effect of irradiation on an ice-rich surface layer and a bluish collisional resurfacing by buried non-irradiated layers (Luu & Jewitt 1996). A strong correlation of color vs. orbital inclination detected among the dynamic family of classical KBOs (Trujillo & Brown 2002) led to a further development of collisional resurfacing models, accounting for differently irradiated subsurface layers (Gil-Hutton 2002), or for collision-triggered cometary activity and size-dependent resurfacing (Delsanti et al. 2004). While these models had difficulties in explaining the detection, or non-detection, of several surface properties and correlations between these and the orbital properties (e.g., Jewitt & Luu 2001; Thébault & Doressoundiram 2003; Delsanti et al. 2004), the dynamical model of Gomes (2003) suggested that high-inclination KBOs formed closer to the Sun migrate afterward into their current orbits. This opened a new window for the possibility of original compositional differences among KBOs (e.g., Tegler et al. 2003).

Simultaneously, a debate on the global existence, or absence, of only two distinct surface color groups of Centaurs and KBOs (Tegler & Romanishin 1998, 2003) led to the conclusion that only Centaurs seemed to be separated into two distinct groups, whereas KBOs presented a somewhat continuous distribution of colors (Peixinho et al. 2003), and current data suggest that the existence of two distinct groups of objects is related to their sizes and not to their dynamical families (Peixinho et al. 2012; Fraser & Brown 2012).

Dynamical models indicate that the giant planets have migrated throughout the history of the solar system, sculpting the structure of the current Kuiper Belt (e.g., Levison et al. 2011; Batygin et al. 2011, and references therein). Although there is still a debate on the details of this process, it became more plausible that the Kuiper Belt is composed by a mixture of bodies formed at distinct heliocentric distances and, therefore, their distinct surface properties would be primarily caused by primordial

compositional differences (e.g., Brown et al. 2011). The most recent analysis of albedos vs. visible colors of Centaurs and KBOs identifies a surface color and albedo separation that is also evidence for such primordial compositional differences (Lacerda et al. 2014). Furthermore, latest laboratory works show that it is possible to reproduce a wide range of surface colors of Centaurs and KBOs with an appropriate combination of initial albedos, collisional evolution, and space weathering, implying that it is unlikely that a straightforward dependence between the dynamical properties and the surface properties exists (Kaňuchová et al. 2012). Thus, it comes as no surprise that the two taxonomy schemes proposed, so far, identify several groups with distinct surface properties (Barucci et al. 2005; Perna et al. 2010; Fraser & Brown 2012; Dalle Ore et al. 2013), failing to detect any clear dependence between the dynamical properties and the surface properties, although the proportions of taxonomic families vary among the dynamical families.

With the growing interest on detailed spectroscopy studies in the near-IR, where several ice absorption bands can be detected, as well as with the existence of an apparently large database of visible colors of our objects, it might be perceived as irrelevant to continue photometric studies of KBOs and their progeny. However, spectroscopic studies are still limited to fewer than 100 objects (i.e., $\sim 5\%$ of known objects) and are extremely biased toward only the brightest objects, whereas visually all detected objects might be studied photometrically, and, as aforementioned, it is currently known that smaller (fainter) objects do possess a peculiar bimodality of surfaces that is not seen among the larger (brighter) objects. The question is, nonetheless, pertinent, as in theory one does not need to study all the population but only a representative sample of it. The problem is that the more subtle the effect one seeks, the larger the sample required.

In this work we exclusively focus on studying the correlations between the visible surface colors and the orbital parameters for each traditional dynamical family of KBOs and Centaurs of our own compilation of visible colors available in the literature, assumed to be unbiased samples of each family, by default. We analyze and discuss for the first time, what we might be missing as a function of the available sample sizes for each dynamical family, and also as a function of the observational error-bars. For an analysis of the (statistical) distributions of the surface properties of each dynamical family see Hainaut et al. (2012).

2. Data sample

Orbital elements were gathered from the Asteroid Orbital Elements Database, *astorb.dat*¹, maintained by the Lowell Observatory based on astrometric observations by the Minor Planet Center. Orbital inclinations i are relative to the ecliptic and not to the so-called Kuiper Belt plane.

We used the classification scheme suggested by Lykawka & Mukai (2007), including their analysis of objects located in the mean motion resonances (MMR) with Neptune, following a ten-step algorithm:

1. in 3 : 2 MMR with Neptune $\Rightarrow$ Plutino
2. in other MMR with Neptune $\Rightarrow$ other resonant
3. $q < a_J \Rightarrow$ not analysed
4. $q > a_J \wedge a < a_N \Rightarrow$ Centaur
5. $a_J < q < a_N \wedge a \geq a_N \Rightarrow$ scattered disk object (SDO)
6. $a_N < q \leq 37 \text{ AU} \Rightarrow$ scattered disk object (SDO)
7. $q \geq 40 \text{ AU} \wedge a \geq 48 \text{ AU} \Rightarrow$ detached KBO (DKBO)

¹ <ftp://ftp.lowell.edu/pub/elgb/astorb.dat.gz>, with epoch 20120420

8. $37 AU \leq q \leq 40 AU \Rightarrow$ scattered or detached KBO (SD-KBO)
9. $i < 5^\circ \wedge \{[q \geq 37 AU \wedge (37 AU \leq a \leq 40 AU)] \vee [q \geq 38 AU \wedge (42 AU \leq a \leq 48 AU)]\} \Rightarrow$ cold classical KBO (cCKBO)
10. $i \geq 5^\circ \wedge q \geq 37 AU \wedge (37 AU \leq a \leq 48 AU) \Rightarrow$ hot classical KBO (hCKBO).

We are aware that there are more complex classification schemes that may be more rigorous, but the boundaries between families do not change significantly. We chose this one for its computational simplicity, which provides a dynamical classification for all objects in our sample, whereas dynamical simulation-based classifications are only available for a limited number of objects (e.g., Gladman et al. 2008).

The so-called Haumea family (Brown et al. 2007; Snodgrass et al. 2010) — (136108) Haumea, (24835) 1995SM₅₅, (55636) 2002TX₃₀₀, (86047) 1999OY₃, (19308) 1996TO₆₆, (120178) 2003OP₃₂, (145453) 2005RR₄₃, 2003SQ₃₁₇, 2003UZ₁₁₇, and 2005CB₇₉ — are extremely peculiar objects, therefore we removed them from the statistical analysis. We also removed the two objects in retrograde orbits (2008KV₄₂, 2008YB₃). For a total of 366 objects in our database, we carried out a global analysis, that is, all objects except Haumea family and retrograde orbits, of $n = 354$ objects, calling it All*; another analysis also removed Centaurs, which we call KBOs*; a third analysis according to dynamical family; and finally, an analysis for which we combined some families.

We compiled the visible colors (*BVRI* broadband filters or equivalent filters transformed to that system) available for 366 objects (see Table 5) and fully trusted their published error-bars. For each input paper and each object or observation, we computed the reflectance spectrum using Eq. (3) from Delsanti et al. (2001), when two or more filters were available. The resulting spectra were manually checked and obviously deviating data from a given filter were removed from the dataset. Color indexes were computed within one given epoch, which led to colors obtained from simultaneous photometry (the different bands were observed over a maximum time-span of two hours). Then the average colors indexes and their one-sigma errors from different papers and epochs were computed for each object using Eqs. (1) and (2) from Hainaut & Delsanti (2002), which provided more accurate estimates when multiple measurements are available. The absolute magnitude in *R* band (H_R) was computed for each object and epoch whenever an *R*-band magnitude was available, using $H_R = R - 5 \log(r\Delta)$, where R is the *R*-band magnitude and r and Δ the helio- and geocentric distances at the time of observations, respectively. Different values for a given object were also averaged using Eqs. (1) and (2) from Hainaut & Delsanti (2002). We did not correct for any phase effect, as already discussed in Peixinho et al. (2012); see Table 5.

3. Statistical methods

3.1. Statistical tests

- **Spearman-rank correlation accounting for data error-bars:** We analyzed the possible existence of correlations between surface colors of Centaurs and KBOs and orbital parameters using the Spearman-rank correlation coefficient ρ (Spearman 1904). The significance of ρ was estimated using $t = \rho[(n-2)/(1-\rho^2)]^{1/2}$, which obeys approximately a Student *t*-distribution with $n-2$ degrees of freedom, where n is the number of data-points (Kendall et al. 1939).

To take into account the effect of observational errors, we considered each data-point not as fixed, that is, exact, but as a Gaussian distribution centered on the measured value with a standard deviation equal to the corresponding error-bar. Then, we extracted 1000 new random data-sets in which each data-point followed its own Gaussian probability distribution. Since the Spearman ρ -distribution is not Gaussian, we used the Fisher transformation $\zeta(\rho) = \frac{1}{2} \ln\left(\frac{1+\rho}{1-\rho}\right) = \arg \tanh(\rho)$ — Fisher (1915) —, determined the mean $\langle \zeta_\rho \rangle$, that is, the most probable ζ_ρ value, the $+\sigma_\zeta/-\sigma_\zeta$, and transformed it back to ρ -values using $\rho_\zeta = \tanh(\zeta_\rho)$, obtaining the most probable correlation value $\langle \rho_\zeta \rangle$, and the shortest interval containing 68.2% of the distribution, which is the equivalent of a Gaussian $\pm 1\sigma$ limits, which in this case are not necessarily symmetrical. We denote this as $+\sigma_e/-\sigma_e$ (see Peixinho et al. 2004, for more details).

This method yields the confidence interval of the most probable ρ of our sample, but does not yield the confidence interval for the correlation value of the parent population. For that we made 1000 bootstrap extractions from our data-points and computed the corresponding $\langle \rho_b \rangle$ and $+\sigma_b/-\sigma_b$, analogously to the previous case (e.g., Efron & Tibshirani 1993).

Usually, the $\pm\sigma_b$ interval dominates over $\pm\sigma_e$, but a more accurate estimation of the confidence interval for the correlation of the parent population is given by quadratically adding $\sqrt{(\sigma_e^2 + \sigma_b^2)}$, which, after reconversion gives us the best estimate for the 68.2% confidence interval: $\langle \rho \rangle_{-\sigma^+}^{+\sigma^+}$.

– Partial correlation

When dealing with multivariate variables, we constantly face the same question: given three variables A, B, and C, is the correlation observed between the two variables A and B (ρ_{AB}) due to the association of A and B with the third variable, C, and not to the direct association between them? Using the partial correlation, we may eliminate the suspected effects of this third variable C on the relation between A and B by making C somehow constant, denoted by $\rho_{AB.C}$. This is achieved by

$$\rho_{AB.C} = \frac{\rho_{AB} - \rho_{AC}\rho_{BC}}{\sqrt{(1-\rho_{AC}^2)(1-\rho_{BC}^2)}}. \quad (1)$$

If $\rho_{AB.C}$ does not differ significantly from ρ_{AB} , then the association between A and B is most probably direct and not masked by their correlations with C. To take into account the effects of data error-bars, we have used the same process as previously described for the Spearman-rank correlation. The significance of $\rho_{AB.C}$ is estimated analogously to the case of ρ , but with one less degree of freedom, that is, using $t = \rho_{AB.C}[(n-3)/(1-\rho_{AB.C}^2)]^{1/2}$ — see, for instance, Macklin (1982).

3.2. Assessing the risk of missing existing correlations

Searching for correlations means testing a null hypothesis (H_0), that is, variables X and Y do not show evidence for correlation, against an alternative hypothesis (H_1), where variables X and Y show evidence for correlation. This evidence is measured by the significance level (SL), confidence level ($CL = 1 - SL$), or *p*-value ($= SL$). The threshold *p*-value below which we reject H_0 is called α . The lower the significance, the lower the probability of incorrectly rejecting the null hypothesis (type I error). To diminish the odds of finding false correlations, we simply adopted a stricter criterion for rejecting H_0 , for instance, instead of accepting an $\alpha = 1\%$ probability of falsely finding a correlation, we

accept only $\alpha = 0.1\%$, or even lower. The drawback is that this increases the chances of not detecting correlations that might in fact exist, meaning that we increase the *risk*, β , of missing some real correlations (type II error), or decrease the $(1 - \beta)$ power to find them. If we wish to diminish this risk, we must either increase the α threshold with which we reject the null hypothesis or increase the sampling. We have analyzed the sample size requirements, n , to detect a correlation at some given significance level (higher or equal to α) and to only miss a β proportion. Since we wish to compare the case where the correlation is indeed $\rho \neq 0$ with the case of $\rho = 0$, from Modarres (1996) and Bonett & Wright (2000) we obtain

$$\sqrt{n(n-3)} = \frac{4(z_{1-\alpha/2} + z_{1-\beta})^2(1 + \rho^2/2)}{(\ln \frac{1-\rho}{1+\rho})^2}, \quad (2)$$

where $z_{1-\alpha/2}$, and $z_{1-\beta}$ are the quantiles of a normal distribution for our threshold $\alpha/2$ — since we are using two-tailed tests — and risk β , respectively, from which we compute the smallest integer not lower than n , that is, $\lceil n \rceil$, ceiling of n . For example, to see evidence for a correlation at a 2.5σ level, that is, $\alpha = 0.0124$ and $z_{1-\alpha/2} = 2.5$, in a sample extracted from a parent population which we knew to possess $\rho = 0.6$ accepting only a $\beta = 0.1$ probability (risk) of missing it, that is, $z_{1-\beta} = 1.2816$, we would need a sample of $n \geq 39$. If we accept a higher risk of missing the detection, for example, $\beta = 0.2$ ($z_{1-\beta} = 0.8416$), we would need only $n \geq 31$.

3.3. Degraded correlations

When taking into account the data error-bars, the possible correlations between the X and Y variables will diminish or degrade. That is, taking each data-point not as an exact value but as a Gaussian probability distribution centered on the measured value diminishes the possible association strength between the two variables. The larger the error-bars, the stronger this effect.

The error-bars of surface colors of Centaurs and KBOs tend to become larger with increasing wavelength (e.g., from visible $B - V$ to the near-infrared $H - K$). It might be possible that two different pairs of color indexes, X_1 vs. Y_1 and X_2 vs. Y_2 , correlate with equal strength (e.g., $B - V$ vs. $V - R$ and $V - R$ vs. $R - I$) — if we precisely knew the full population, for instance. However, not only are we estimating the populations' ρ s from a limited sample, but if the error-bars of (X_2, Y_2) are larger than those of (X_1, Y_1) we might also obtain a $|\rho_2| < |\rho_1|$ or even lose evidence for correlation on (X_2, Y_2) . To analyze this possibility, we performed the following study:

1. We generated a parent population of 1000 (X, Y) pairs with a desired correlation value. Each variable follows a Gaussian distribution, and their values are renormalized into the interval $[0, 1]$ (the mean will become 0.5) by two distinct methods: a) the lowest value will correspond to zero, the highest to one; b) the mean value minus 3σ will correspond to zero, the mean value plus 3σ to one (with this renormalization some points may fall outside the $[0, 1]$ interval).
2. We extracted from that parent population a sample of n elements. We added Gaussian error bars to the Y elements and generated 1000 new samples from the n elements by the same process as we used for the real data-samples (see Sect. 3.1). The approximate sampling error given by Bonett & Wright (2000),

$$\sigma_{\rho}^2 \approx (1 + \rho^2/2)/(n - 3), \quad (3)$$

was added quadratically to the dispersion of the Fisher-transformed simulations — instead of using bootstraps extractions — before reconvert them to obtain a final $\langle \rho \rangle_{-\sigma}^{+\sigma}$.

3. We iterated the two previous steps for $\rho = 0.8, 0.7, 0.6$, and 0.5 , each with samplings of $n = 15, 20, 30, 50, 75$, and 100 , and varied the observational error-bars from $\sigma_{obs} \equiv \sigma_o = 0.00$ to 0.30 in steps of 0.02 .
4. We repeated these cycles with X and Y following a random uniform distribution instead of a Gaussian.

First, the simulations show that drawing the sample from a Gaussian distribution or from a uniform distribution, either renormalized into the interval $[0, 1]$ by the methods 1a) or 1b), has little to no influence on the degradation of ρ values and their error-bars, with increasing simulated observational errors σ_o . Variations are on the order of 0.01 or lower. Hereafter, the analysis refers to the simulations drawn from a Gaussian distribution that was renormalized using method 1a).

Second, we find that the effects of the (simulated) observational errors on X and on Y , which we denote by σ_{oX} and σ_{oY} , are equivalent to performing the simulations with additional observational errors only on the X variable of $\sigma_o = \sigma_{oX} + \sigma_{oY}$, adding no errors to the Y variable. For example, having a series of both X and Y values with error-bars of 0.1 has an equivalent effect on the correlation values as having a series of X values with error-bars of 0.2 and Y values with no errors. This result is most convenient and simplifies the rest of our analysis.

Third, we find, empirically, that for each initial ρ value its degradation with increasing observational errors σ_o is independent of the sample size n . The estimated 68.2% error interval on the measured $\langle \rho \rangle_{-\sigma}^{+\sigma}$ changes with n because it is largely dependent on it (see Eq. 3), but the diminishing of $\langle \rho \rangle$ shows no evident dependence on n .

Using a Levenberg-Marquardt algorithm (Levenberg 1944; Marquardt 1963), we find that third-degree polynomials fitted with great accuracy (all cases with residuals $rms < 0.0063$) the detectable ρ functions denoted as $\rho^* = f(\rho, \sigma_o)$:

$$\rho^* = \begin{cases} 1.000 - 0.950\sigma_o - 10.544\sigma_o^2 + 25.349\sigma_o^3 & : \rho = 1.0 \\ 0.900 - 0.497\sigma_o - 8.668\sigma_o^2 + 18.387\sigma_o^3 & : \rho = 0.9 \\ 0.800 - 0.309\sigma_o - 7.255\sigma_o^2 + 14.435\sigma_o^3 & : \rho = 0.8 \\ 0.700 - 0.176\sigma_o - 6.373\sigma_o^2 + 12.299\sigma_o^3 & : \rho = 0.7 \\ 0.600 - 0.129\sigma_o - 4.322\sigma_o^2 + 7.461\sigma_o^3 & : \rho = 0.6 \\ 0.500 - 0.0513\sigma_o - 2.817\sigma_o^2 + 4.373\sigma_o^3 & : \rho = 0.5 \end{cases} \quad (4)$$

An immediate consequence of the ρ degradation with larger σ_o s is the magnitude decrease of the detectable ρ^* . That is, even if we had a parent population such that $X = Y$, hence $\rho = 1$, the fact that a sample has error-bars will impose a limit on the detectable correlation, regardless of the sample size n . In practice, from the correlation analysis point of view, this means that it is very important to diminish the observational errors, but rather pointless to enlarge the sampling with the same or larger error-bars if the elements in the sample are already representative of the whole population, of course, and the significance level of the detection has reached the threshold we desire.

Another consequence is that increasing β increases the risk of missing an existing correlation. Therefore, to keep β constant (for a theoretically detectable correlation), we need to increase the sampling n . We return to the example in Sect. 3.2. We know

the parent population has $\rho = 0.6$ and we need $n \geq 31$ to detect evidence for a correlation at a 2.5σ level with a risk $\beta = 0.2$ of missing it. Suppose now that each X data-point has an observational error $\sigma_{oX}=0.1$ and each Y data-point also has $\sigma_{oY}=0.1$. As we saw earlier, this is equivalent to transferring all the error-bars to the X variable, such that $\sigma_o = \sigma_{oX} + \sigma_{oY} = 0.2$, and to assume Y with no errors. (Note that we take X and Y as normalized to $[0, 1]$ so this error represents 10% of the full range of values on X and 10% on Y .) From Eq. 4 we see that with such observational errors the detectable correlation drops to $\rho^* = 0.46$, and now, from Eq. 2, to detect evidence for a correlation at the same 2.5σ significance level, we will need $n \geq 53$ data-points, that is, 70% more data-points ($n \geq 67$ for $\beta = 0.1$). In this work, we are able to quantitatively predict the sample size needed for the cases studied below.

3.4. Correction for the false-discovery rate of the correlations

Since for each sample or subsample we performed about $m = 150$ correlation tests, the chances of finding evidence for a correlation increase, even if there is no correlation at all (see Miller 1981, for a review). For example, suppose we reject the null hypothesis of no correlation using a threshold $\alpha = 0.0124$ (2.5σ level). There is a 1.24% probability of obtaining evidence for a correlation just by chance, even if it is not there. This means that performing 150 tests will probably lead to one or two false discoveries. To solve this problem, we used the false-discovery rate (FDR) procedure as proposed by Benjamini & Hochberg (1995). That is, instead of ensuring that the probability of erroneously rejecting even one of the null hypothesis is always lower than a certain α — using the Bonferroni procedure of rejecting H_0 only when its p-value is $\leq \alpha/m$ (Bonferroni 1935)—, we ensure that the proportion of incorrectly rejected H_0 's, or false discoveries, is α .

The Benjamini-Hochberg procedure works as follows:

1. Order the m p-values from all tests as $p_1 \leq p_2 \leq \dots \leq p_m$;
2. find the largest integer k such that $k = \max\{i : p_i \leq \frac{i}{m}\alpha\}$,
3. reject all the null hypotheses whose p-value is $\leq p_k$.

This procedure ensures that $\text{FDR} \leq \alpha$. That is, if we use $\alpha = 0.0124$ (2.5σ) at most 1.24% of all the positive results (i.e., all the rejected null hypotheses) may be false positives. For example, if we simultaneously test 150 hypotheses and reject fewer than 30 of them, or in other words, we have fewer than 30 discoveries, it is unlikely that we make even one false discovery (at 2.5σ , naturally).

4. Reddening line

Delsanti et al. (2001) introduced the concept of drawing a line in the color-color plots indicating the location of objects that probably possess linear reflectivity spectra. As discussed in detail later by Delsanti et al. (2004), a linear reflectivity spectrum does not result in a linear relation between colors, and vice versa. This line has been widely used to infer the linearity or non-linearity of object spectra. However, each telescope has its own set of filters, and the differences between the effective central wavelength for the same filters among the different sets has a significant effect on the location of this reddening line. This effect can be strong for the visible BVR colors, but it is not very significant for the VRI bands.

We have computed the reddening line for several of the most frequently used telescopes or instruments for KBO observations

Table 1. Central wavelengths in Å of different telescope filters

Telescope	Filters			
	B	V	R	I
8.2m ESO-VLT	4237	5481	6480	7935
3.6m ESO-NTT	4212	5442	6416	7950
10m Keck	4377	5473	6417	7599
8.2m Subaru	4400	5500	6600	8050
3.6m CFHT	4312	5374	6581	8223
Average	4307.6	5454	6498.8	8016.2

and analyzed their differences. As seen in Fig. 1, the reddening line obtained for the Mould- BVR filters of the CFHT12k camera of the 3.6m CFHT telescope is higher, and the reddening lines obtained for the ESO FORS1 camera of the 8.2m VLT telescope or the ESO SuSi2 camera of the 3.6m NTT telescope are the lowest. The redenings for the LRIS camera on the 10m Keck telescope and the FOCAS cameras on the 8.2m Subaru telescope are intermediate. Therefore, interpretations based on a precise location of this line should be taken with caution. We have used the solar colors $(B - V)_\odot = 0.65$, $(V - R)_\odot = 0.36$, and $(V - I)_\odot = 0.70$ from Ramírez et al. (2012). Small changes in the solar colors do not affect the reddening line.

5. Data analysis

This work is focused on $BVRI$ surface information or properties of Centaurs and KBOs. We have analyzed the correlations between the color indexes $B - V$, $V - R$, $R - I$, $V - I$, $B - I$, and $B - R$, and the spectral gradient, Grt — also known as the slope parameter or reddening S , which expresses the reflectivity spectrum variation in percent of reddening per 1000 Å (Hainaut & Delsanti 2002) —, and the correlations between these indexes and the absolute R -magnitude not corrected for phase effects, that is, $H_R(\alpha) \equiv R(1, 1, \alpha)$. Correlations between these properties and the orbital parameters perihelion, q , aphelion, Q , orbital inclination relative to the ecliptic, i , orbital eccentricity, e , and semi-major axis, a , have also been studied. For some families we have also analyzed the correlation with the Tisserand parameter relative to Neptune, T_N , the Tisserand parameter relative to Jupiter, T_J , the mean random impact speed velocity, v_{rms} , the orbital excitation, ε , and Öpik's ψ . These parameters are explained below when used. As mentioned in Sect. 2, we relied entirely on the published error bars, although for multiple measurements we checked for deviant points before carrying out the weighted averaging. Possible over- or underestimated errors for some data-points, most probable in the case of single measurements, are not expected to have any statistical influence on the methods we used and developed, unless if generalized.

Each dynamical family, or group of families, was subject to 84 tests for correlation. The exceptions were i) Centaurs, which were subjected to 92 tests due to the inclusion of T_J in the tests; ii) classical KBOs, which were subjected to 100 tests, given the inclusion of v_{rms} , ε , and ψ ; and iii) resonant objects were only subjected to 76 tests, since correlations with T_N were not tested. These numbers are extremely important to correct for the false-discovery rate (FDR) as described in Sect. 3.3.

We discuss the correlations for each dynamical family separately from Sects. 5.1 to 5.6. KBOs, as a whole, are discussed in Sect. 5.7, binary or multiple objects in Sect. 5.8, and all objects




Fig. 1. $(B - V)$ vs. $(V - R)$ — top — and $(V - R)$ vs. $(R - I)$ — bottom — reddening lines obtained with the most frequently used sets of BVR filters in our database and the computed average reddening line. For the $(B - V)$ vs. $(V - R)$ case different filters have a strong effect on the location of the reddening line. For the $(V - R)$ vs. $(R - I)$ case the effect is almost negligible.

— excluding Haumea family and retrogrades — are discussed in Sect. 5.9. For each case we also comment on the sample size and the consequent risk of missing correlations. Strictly speaking, this discussion is valid for a characterization of these families and groups under the assumption that the data are an unbiased sample of the parent population. The statistical tools, however, are general and can be applied to other classification systems.

5.1. Centaurs

Our sample has $n = 33$ Centaurs. Because one of them was a retrograde object, however, we removed it and analyzed a total of $n = 32$ Centaurs. All of their surface colors indexes strongly correlate with each other, which indicates a strong degree of predictability of one color when knowing another (see middle panels of Fig. 2 and Table 2). No correlation between colors and absolute magnitude, and between these and any orbital parameter reaches a 2.5σ level after correcting for FDR. We stress that with $n = 32$ objects, the lowest correlation value we expect not

to miss at a 10% risk β and at 2.5σ is $\rho_{\beta 2.5\sigma} = 0.64$. At the same 10% risk, but at 3σ confidence level we have $\rho_{\beta 3\sigma} = 0.70$. In other words, with the current sampling of Centaurs there is a risk higher than 10% of missing any possible strong correlation. Increasing the sampling of Centaurs is still very important. To have a risk lower than 10% of missing any correlation stronger than ≈ 0.5 , we need a sample of $n \geq 70$ objects (more specifically, $\rho_{\beta 2.5\sigma} \leq 0.46$ and $\rho_{\beta 3\sigma} \leq 0.50$).

5.2. Scattered disk objects

The dynamical boundaries for the definition of scattered disk objects, or SDOs, are ill defined. Different authors used slightly different definitions. Following the definition we used, we have $n = 62$ SDOs. Since one of them is a retrograde object and another one a Haumea family object, we have analyzed a total of $n = 60$ SDOs. Most color indexes correlate with each other (see bottom panels of Fig. 2 and Table 2). The exception are $(B - R)$ vs. $(R - I)$ and $(V - R)$ vs. $(R - I)$, which indicates that it is difficult to extrapolate color values, although interpolating them seems possible. There is a correlation between H_R and q that we interpret as a biasing effect among SDOs, meaning that for orbits with high perihelia only the brightest objects can be detected, which was also discussed by Santos-Sanz et al. (2012) — for a review on biasing effects see Kavelaars et al. (2008). At $\beta = 0.1$, with $n = 61$, we have $\rho_{\beta 2.5\sigma} = 0.48$ and $\rho_{\beta 3\sigma} = 0.53$, so some medium strength correlations might still be missed. A $\sim 50\%$ increase in the sampling of SDOs, to $n = 90$, would lower these values to $\rho_{\beta 2.5\sigma} = 0.39$ and $\rho_{\beta 3\sigma} = 0.44$. Note, however, that this biasing will not be removed solely by increasing the sample.

5.3. Scattered or detached Kuiper Belt objects

According to the classification scheme we have used, our sample has $n = 7$ detached KBOs (one of which is a binary object) and $n = 15$ scattered or detached KBOs (three of which are binaries). We opted to analyze them as one group (SDKBOs). The data show strong correlations between colors that we may consider as interpolations if we convert them into a spectrophotometric spectra. To illustrate, if we have a flat spectrum, the colors $(B - R)$ or $(V - I)$, for example, can be seen as mere interpolations of the $(B - I)$ color. Analogously, when in presence of a flat spectrum, the $(B - I)$ color, for example, could be obtained by extrapolating from $(B - R)$. This means that similarly as for the SDOs, consecutive colors or color extrapolations do not show evidence of correlation (see bottom panels of Fig. 2 and Table 2). For example, $(B - I)$ vs. $(B - R)$ are strongly correlated, but the wavelength $(B - R)$ is inside the $(B - I)$ wavelength range, while $(B - R)$ vs. $(R - I)$ are not correlated. Because we are dealing with small-sampling values the $\beta = 0.1$ risk shows that it is easy to miss even very strong correlations among SDKBOs. For $n = 20$, $\rho_{\beta 2.5\sigma} = 0.76$ and $\rho_{\beta 3\sigma} = 0.82$. Clearly, it would be good to triple the sample.

5.4. Plutinos

Our sample has $n = 49$ Plutinos, or $(3 : 2)$ resonants, including Pluto and four binaries. All color indexes correlate with each other except for $(V - R)$ vs. $(R - I)$ (see Fig. 3 and Table 2). Nonetheless, $(B - V)$ correlated with $(V - R)$, and $(B - R)$ correlates with $(V - I)$. Therefore, either color interpolation or extrapolations might be possible, but for the latter, B -band information would be required. No correlations between colors and




Fig. 2. $(B - V)$ vs. $(V - R)$ plots — left column — and $(V - R)$ vs. $(R - I)$ plots — right column — including the reddening line. Solar colors are indicated with the $\odot$ symbol. Top panels: All objects, Haumea family objects, and retrograde objects. Middle panels: Centaurs. Bottom panels: Scattered disk objects (SDOs), and scattered or detached Kuiper Belt objects (SDKBOs). The stronger correlation between $(B - V)$ and $(V - R)$ colors than between $(V - R)$ and $(R - I)$ is clear. Apart from the solar-like colors of Haumea family objects and the two color groups of Centaurs in the $(B - V)$ vs. $(V - R)$ there is no apparent relevant difference between the color distribution of each family of objects.

orbital parameters reach a 2.5σ level after correcting for FDR. Given the non-detection of the $(V-R)$ vs. $(R-I)$ correlation with $n = 42$ objects, we highlight that at $\beta = 0.1$ risk $\rho_{\beta 2.5\sigma} = 0.56$ and $\rho_{\beta 3\sigma} = 0.63$. As for the SDOs, some medium-strength correlation might be missed here. Increasing the sample by $\sim 50\%$ is recommended.

5.5. Other resonants

Our sample has $n = 73$ non-Plutino resonants: $n = 2$ (1 : 1), or Neptune Trojans; $n = 3$ (5 : 4); $n = 6$ (4 : 3); $n = 1$ (11 : 8); $n = 1$ (18 : 11); $n = 10$ (5 : 3) — 1 binary —; $n = 1$ (12 : 7) — Haumea multiple system —; $n = 1$ (19 : 11) — Haumea family member —; $n = 15$ (7 : 4) — 1 binary —; $n = 1$ (9 : 5); $n = 10$ (2 : 1) — 3 binaries —; $n = 3$ (9 : 4); $n = 3$ (7 : 3); $n = 2$ (12 : 5); $n = 12$ (5 : 2) — 1 binary —; $n = 1$ (8 : 3) — binary —; $n = 2$ (3 : 1); $n = 1$ (11 : 2); and $n = 1$ (11 : 3). We have removed the Haumea family members from the sample — (136108) Haumea and (019308) 1996TO₆₆ — because of their extremely peculiar behaviors. This means that we analyzed the correlations for $n = 71$ objects (see Fig. 3 and Table 2). Absolutely all color indexes correlate with each other at a 3σ confidence level after the FDR correction. The correlation magnitude weakens to $\rho < 0.6$ only for $(B-R)$ vs. $(R-I)$, $(B-V)$ vs. $(R-I)$, and $(V-R)$ vs. $(R-I)$. There is a moderate correlation between H_R and q , suggesting that we have an absolute magnitude bias among these objects similarly as for the SDOs. As for the previous cases, we detect no correlations between color indexes and orbital parameters. For $n = 71$, the sample size at which the lowest ρ is detected, the $\beta = 0.1$ risk implies $\rho_{\beta 2.5\sigma} = 0.44$ and $\rho_{\beta 3\sigma} = 0.50$. Therefore, only weak or moderate correlations might be missed. As a whole, the other resonants do not require a larger sampling to detect any high relevant correlation. However, it is not likely that all the different resonances will possess the same surface color behaviors, and each resonance should be sampled such that they can be analyzed independently of the others. This still requires a large effort since even with $n = 20$ objects one will only be able to be 90% certain to detect very strong correlations ($\rho_{\beta 2.5\sigma} = 0.76$ and $\rho_{\beta 3\sigma} = 0.81$).

5.6. Classical KBOs

Classical KBOs, also known as classicals or CKBOs, are known for their color-inclination correlation, which is not clearly detected in any other family of KBOs (see Fig. 4 top). As for the case of other resonants, we removed the seven Haumea family objects from the classicals sample: 2003SQ₃₁₇, 2003UZ₁₁₇, 2005CB₇₉, (024835) 1995SM₅₅, (055636) 2002TX₃₀₀, (120178) 2003OP₃₂, and (145453) 2005RR₄₃. When plotting $B-R$ vs. orbital inclination i , we have three outliers with low $B-R$ colors at very low inclination (see also Peixinho et al. 2008). Romanishin et al. (2010) analyzed a subfamily of classical objects with semi major-axes below the nominal location of the (3 : 2) resonance, calling them inner classicals. That subfamily presented a bimodal behavior of surface colors, unlike the rest of classical objects, and probably a distinct origin from the primordial protoplanetary disk. Although we did not use the same classification scheme, the outliers 1998WV₂₄, and 2003YL₁₇₉ belong to that group, and we decided to remove them from the analysis. There is, however, no similar indication why object 2002VD₁₃₁ might be an outlier, but because of its extreme outlier behavior

we chose to eliminate it as well. The final sample we analyzed has $n = 117$ classical objects (see Table 2).

5.6.1. Correlation of color vs. color of classical KBOs

CKBOs evidence neither $(B-V)$ vs. $(V-R)$ nor $(V-R)$ vs. $(R-I)$ correlations. And with their sample size, anything stronger than $\rho \sim 0.43$ should be detected at least at a 2.5σ level, although within the ρ error-bars we see that these correlations among KBOs*, taken globally, increase when CKBOs are removed from the sample (see Table 2). The middle panel of Fig. 3 shows that this result seems clear. Interestingly, there are more CKBOs in the $(B-V)$ vs. $(V-R)$ plot far below the reddening line than for any other family. We return to this question in Sect. 5.9.2.

5.6.2. Correlation of color vs. orbital parameters of classical KBOs

The trend between the colors of classical objects and their orbital inclination was first reported by Tegler & Romanishin (2000). Explicit computations of a correlation were made later by Trujillo & Brown (2002) and Hainaut & Delsanti (2002).

This correlation was thought to be caused by some collisional evolution mechanism, known as collisional resurfacing (CR) — see Luu & Jewitt (1996) — which seemed highly dependent on the orbital inclination because of its link with a stronger or weaker collisional environment. Different parameters have been used to analyze it, such as collisional velocity, orbital excitation, ψ parameter, or simply orbital inclination. We analyzed each one of them, also including the Tisserand parameter relative to Neptune (T_N).

Collisional velocity v_c : The mean square of a KBO encounter velocity with its circular analog, that is, a non-inclined circular orbit with heliocentric distance a , is given by

$$v_{rms} = \sqrt{\frac{GM_{\odot}}{a}(e^2 + i^2)} = v_K \sqrt{e^2 + i^2}, \quad (5)$$

where e represents the orbital eccentricity, i stands for the orbital inclination relative to the ecliptic, and a for the semi-major axis. If a is measured in AU, then $v_K \approx 29.8/\sqrt{a}$ [$km\ s^{-1}$]. For simplicity, we call this v_{rms} simply as collisional velocity v_c .

Trujillo & Brown (2002) discussed the possible link between the color-inclination correlation and impact velocity but, given the lack of detailed modeling available at the time, they also discussed the possibility that the correlation might be a consequence of multiple subpopulations with different primordial colors and inclinations. Stern (2002) explored the link between KBO colors and their mean collisional velocities in more detail and also accounted for the gravitational effect of each object using $v_{coll} = \sqrt{v_{esc}^2 + v_{rms}^2}$, where v_{esc} is the KBO's escape velocity, calculated assuming a bulk density $\rho = 1500\ kg\ m^{-3}$ and a geometric albedo $p = 0.04$. Stern's Spearman-rank correlations of $(B-R)$ vs. v_{coll} and $(B-V)$ vs. v_{coll} were $\rho_{BR v_{coll}} = -0.39$ ($p = 0.005$) and $\rho_{BV v_{coll}} = -0.44$ ($p = 0.0005$), respectively, for a total of $n = 81$ KBOs. Note that these correlation values are weaker than those reported by Trujillo & Brown for colors vs. inclination of classical KBOs also because Stern analyzed all families together. Nonetheless, these works seemed to confirm the existence of a physically meaningful correlation between KBO colors and their mean collision velocities, making a strong case for




Fig. 3. $(B - V)$ vs. $(V - R)$ plots — left column — and $(V - R)$ vs. $(R - I)$ plots — right column — including the reddening line. Solar colors are indicated with the $\odot$ symbol. Top panels: Plutinos and other resonants. Almost all non-Plutino resonants seem to lie on the reddening line in $(B - V)$ vs. $(V - R)$, but their stronger correlation value is within the error-bars of the correlations for other families. Middle panels: Classical KBOs and all other KBOs, except Haumea family and retrogrades. In contrast to all other families, CKBOs do not exhibit a $(B - V)$ vs. $(V - R)$ correlation. The existence of many CKBOs far below the reddening line (in BVR), which indicates the presence of many concave visible spectra, suggests that they may exhibit absorption features in the visible that only detailed spectroscopic studies could verify. Bottom panels: Binary or multiple KBOs.

the CR scenario. From analyzing $n = 50$ CKBOs, Doressoundiram et al. (2002) reported $\rho_{BR v_{rms}} = -0.49$ ($p = 0.0004$). Note that in spite of Stern's care in also accounting for v_{esc} to the collisional velocity from our sample, we measure a median difference between v_{coll} and v_{rms} of only 0.02 km s^{-1} , whereas the median v_{rms} is 0.55 km s^{-1} . This is a very small difference given the uncertainty on the albedos and densities, and is the reason why $v_{rms} \equiv v_c$ was used more often. It is noteworthy that Jewitt & Luu (2001), the proponents of the CR scenario, argued against it as the primary cause of color differences, since KBOs should then also exhibit large color variations with rotation, and observations indicated otherwise. Thébault & Doressoundiram (2003) and Thébault (2003) found additional problems with this scenario. Through a numerical analysis, they computed the kinetic energy KBOs would receive by collisions (hereafter K_{coll}), considering several different swarms of impactors and also different KBO distributions. Results showed correlations between K_{coll} and perihelion, but also that K_{coll} correlated systematically much better with eccentricity than with inclination. Observationally, there was evidence for color vs. perihelion and color vs. inclination correlations but, to date, no relevant correlations between color and eccentricity has been found. Furthermore, from these simulations, K_{coll} of Plutinos (3:2 resonants) was higher than the correlation of any other family, but observationally, their colors did not seem to differ. It is, nonetheless, true that this kind of numerical analysis has not been reanalyzed including the current understanding of the sculpting of the Kuiper Belt through migration processes.

Orbital excitation ε : Hainaut & Delsanti (2002) analyzed the color correlations with the object orbital excitation ε , defined as

$$\varepsilon = \sqrt{e^2 + \sin^2 i}. \quad (6)$$

This ε parameter is related with v_{rms} (see Eq. 5), which is another estimate of the KBO velocity with respect to its circular analog. Using Pearson's correlation coefficient (r_s) instead of Spearman's, they obtained for $(B - R)$ vs. ε of $n=13$ CKBOs $r_s = -0.77$ ($p = 0.002^2$), but no relevant values for any other families ($r_s < -0.1$), nor for KBOs and Centaurs taken together ($r_s = -0.21$, $n=34$).

Öpik's ψ : Santos-Sanz et al. (2009) analyzed the color correlations with a ψ parameter, taken from the theory of Öpik (1976) of close encounters of a small body with a planet, given by

$$\psi = \frac{\frac{5}{8} e^2 + \sin^2 i}{a}. \quad (7)$$

In our context, ψ relates with the average energy of the collisions experienced by a KBO. Again, the similarity between this parameter and the two previous ones is obvious. There is a subtle difference: $\sqrt{GM_\odot \psi}$ is an approximation for the KBO average encounter velocity with targets in circular orbits at the KBO instantaneous heliocentric distance on its elliptical orbit, that is, all local circular analogs. The two previous parameters consider only the impact with the global circular analog. Strangely, Santos-Sanz et al. (2009) reported for $(B - R)$ vs. ψ of $n=64$ CKBOs $\rho = -0.02$. In our analysis we obtain $\rho_{(B-R)\psi} = -0.54^{+0.12}_{-0.10}$ ($p < 10^{-6}$), as roughly expected given all the previous works.

Tisserand parameter T_N : The Tisserand parameter (Tisserand 1896), or invariant, has proven to be a useful tool in comet taxonomy (Kresák 1972; Levison 1996, and references therein). It is a dynamical quantity kept approximately constant during a close encounter interaction between a planet and a minor body — for instance, a KBO could have its orbital parameters changed by a close interaction with Neptune, and yet the Tisserand parameter value would not change, allowing us to keep track of the same object. Since KBOs are mostly under gravitational influence of Neptune, we compute it relative to this planet using

$$T_N = \frac{a_N}{a} + 2 \sqrt{\frac{a}{a_N} (1 - e^2)} \cos i, \quad (8)$$

where $a_N = 30.07 \text{ AU}$ is the semi-axis of Neptune and a , e , and i are the orbital parameters of the KBO. A minor body on a circular orbit, coplanar with Neptune, with $a = a_N$ would have $T_N = 3$. A KBO, being a trans-Neptunian object, will have $a > a_N$. If $T_N > 3$, then it will not cross the orbit of Neptune, whereas if $T_N < 3$ means, generally, a Neptune-crosser. Usage of T_N in KBO taxonomy has been suggested by Elliot et al. (2005). Problems with the interpretation of T_N values arise when high inclination values lead to a $T_N < 3$, suggesting Neptune-crossing, even for impossible cases. For example, a KBO on a circular orbit with $a = 37 \text{ AU}$ and $i = 30^\circ$ would have $T_N = 2.7$ and it would not cross the orbit of Neptune. Note that T_N parameter also relates with the relative encounter velocity between a KBO and Neptune: $v_c \propto \sqrt{3 - T_N}$.

Regardless of the dynamical implications, Jewitt et al. (2007) reported for a total of $n = 15$ CKBOs $(B - R)$ vs. T_N and $(U - B)$ vs. T_N correlation values of $\rho_{(B-R)T_N} = -0.75^{+0.13}_{-0.20}$ ($p = 0.0014$), and $\rho_{(U-B)T_N} = -0.86^{+0.08}_{-0.14}$ ($p = 0.00004$), respectively.

Is there only one correlation? CKBOs show moderate to strong correlations, all of similar magnitudes, both with i , v_c , ε , ψ , and T_N , and a moderate to weak correlation with q . The confidence level for the correlation between color and eccentricity does not reach the 2.5σ threshold after correcting for FDR, and no relevant correlation is detected between color and semi-major axis (see Table 3). From the correlations $\rho_{\psi v_c} = 1.00^{+0.00}_{-0.01}$, $\rho_{\psi \varepsilon} = 0.99^{+0.01}_{-0.01}$, and $\rho_{\varepsilon v_c} = 1.00^{+0.00}_{-0.01}$ it is clear that both ψ , ε , and v_c will be correlated with any other variable with the same strength. Consequently, analyzing for correlations between colors and collisional velocity v_c , or orbital excitation ε , or the Öpik ψ is equivalent and, from a statistical point of view, there is no reason to chose one over the other. Hereafter, we chose to work with v_c since it has been used more often in other works.

Now, we consider the correlations between $(B - R)$, T_N , v_c , i , and q more closely. Given the mutual dependence of most of these parameters, we have carried out a partial correlation analysis to assess whether one of these correlation is dominant and the others are only spurious effects with no physical relevance, but are present because of their dependence on the dominant one (see Sect. 3.1).

The partial correlations between $(B - R)$, q , and i are

$$\begin{cases} \rho_{(B-R)q} = -0.41^{+0.15}_{-0.13} & (p = 0.00026; CL = 3.42 \sigma) \\ \rho_{(B-R)i} = 0.14^{+0.14}_{-0.15} & (p = 0.22; CL = 1.23 \sigma) \end{cases}, \quad (9)$$

showing that the correlation between $(B - R)$ colors and orbital inclination is still present and significant after removing the effect of the link between inclination and perihelion. Its strength,

² P-value is our estimate

Table 3. Spearman ρ correlation matrix between $(B - R)$ and orbital parameters for $n = 79$ CKBOs.

	$B - R$	T_N	ψ	v_c	ε	i	q	e	a
$B - R$	•	$0.50^{+0.11}_{-0.13}$ ($p = 3 \cdot 10^{-6}$)	$-0.54^{+0.12}_{-0.10}$ ($p < 10^{-6}$)	$-0.54^{+0.12}_{-0.10}$ ($p < 10^{-6}$)	$-0.53^{+0.12}_{-0.10}$ ($p < 10^{-6}$)	$-0.50^{+0.13}_{-0.11}$ ($p = 2 \cdot 10^{-6}$)	$0.36^{+0.12}_{-0.13}$ ($p = 0.0011$)	$-0.30^{+0.13}_{-0.12}$ ($p = 0.0078$)	$-0.11^{+0.14}_{-0.14}$ ($p = 0.31$)
T_N		•	$-0.78^{+0.07}_{-0.05}$ ($p < 10^{-6}$)	$-0.77^{+0.07}_{-0.05}$ ($p < 10^{-6}$)	$-0.76^{+0.07}_{-0.06}$ ($p < 10^{-6}$)	$-0.85^{+0.05}_{-0.04}$ ($p < 10^{-6}$)	$0.56^{+0.09}_{-0.10}$ ($p < 10^{-6}$)	$-0.29^{+0.11}_{-0.10}$ ($p = 0.00083$)	$0.20^{+0.13}_{-0.14}$ ($p = 0.026$)
ψ			•	$1.00^{+0.00}_{-0.01}$ ($p < 10^{-6}$)	$0.99^{+0.01}_{-0.01}$ ($p < 10^{-6}$)	$0.93^{+0.02}_{-0.03}$ ($p < 10^{-6}$)	$-0.66^{+0.08}_{-0.07}$ ($p < 10^{-6}$)	$0.68^{+0.07}_{-0.08}$ ($p < 10^{-6}$)	$0.26^{+0.12}_{-0.13}$ ($p = 0.034$)
v_c				•	$1.00^{+0.00}_{-0.01}$ ($p < 10^{-6}$)	$0.91^{+0.03}_{-0.03}$ ($p < 10^{-6}$)	$-0.68^{+0.09}_{-0.07}$ ($p < 10^{-6}$)	$0.71^{+0.06}_{-0.07}$ ($p < 10^{-6}$)	$0.28^{+0.11}_{-0.12}$ ($p = 0.0015$)
ε					•	$0.91^{+0.03}_{-0.04}$ ($p < 10^{-6}$)	$-0.67^{+0.09}_{-0.07}$ ($p < 10^{-6}$)	$0.72^{+0.06}_{-0.07}$ ($p < 10^{-6}$)	$0.30^{+0.11}_{-0.12}$ ($p = 0.00052$)
i						•	$-0.51^{+0.09}_{-0.08}$ ($p < 10^{-6}$)	$0.47^{+0.08}_{-0.09}$ ($p < 10^{-6}$)	$0.13^{+0.12}_{-0.12}$ ($p = 0.13$)
q							•	$-0.70^{+0.07}_{-0.06}$ ($p < 10^{-6}$)	$0.04^{+0.12}_{-0.12}$ ($p = 0.69$)
e								•	$0.63^{+0.07}_{-0.09}$ ($p < 10^{-6}$)
a									•

Notes. Correlation values represented as $\rho^{+\sigma}_{-\sigma}$. Significance p-values are two-tailed.

however, diminishes to a more moderate value. Perihelion is not a function of inclination, and the fact that q and i are correlated could be due to some dynamical evolution mechanism in the way the Kuiper Belt was sculpted, or it may simply be the result of a sampling bias.

The partial correlations between $(B - R)$, T_N , and q are

$$\begin{cases} \rho_{(B-R)T_N,q} = 0.39^{+0.13}_{-0.15} & (p = 0.00035; CL = 3.57 \sigma) \\ \rho_{(B-R)q,T_N} = 0.11^{+0.14}_{-0.14} & (p = 0.35; CL = 0.93 \sigma) \end{cases}, \quad (10)$$

showing that the correlation between $(B - R)$ colors and Tisserand parameter is also still present and significant, although diminished to a more moderate value, after removing the effect of the link between T_N and perihelion, analogously to the previous case.

The partial correlations between $(B - R)$, T_N , and i are

$$\begin{cases} \rho_{(B-R)T_N,i} = 0.17^{+0.15}_{-0.15} & (p = 0.14; CL = 1.47 \sigma) \\ \rho_{(B-R)i,T_N} = -0.17^{+0.15}_{-0.14} & (p = 0.12; CL = 1.54 \sigma) \end{cases}, \quad (11)$$

which shows that none remains after eliminating the effect of the other. This should not be seen as strange given that by definition T_N is highly dependent on i , presenting a $\rho_{T_N,i} = -0.85$, so it is very hard to separate one from the other.

The partial correlations between $(B - R)$, i , and v_c are

$$\begin{cases} \rho_{(B-R)i,v_c} = -0.06^{+0.15}_{-0.15} & (p = 0.59; CL = 0.53 \sigma) \\ \rho_{(B-R)v_c,i} = -0.20^{+0.14}_{-0.13} & (p = 0.074; CL = 1.78 \sigma) \end{cases}, \quad (12)$$

showing also that none remains after eliminating the effect of the other, even if the effect is stronger on inclination without the

influence of collisional velocity than on v_c without the effect of i . Analogously to the previous case, by definition the collisional velocity depends on inclination ($\rho_{v_c,i} = -0.91$), so it is hard to separate them.

Finally, the partial correlations between $(B - R)$, T_N , and v_c are

$$\begin{cases} \rho_{(B-R)v_c,T_N} = -0.27^{+0.14}_{-0.13} & (p = 0.016; CL = 2.41 \sigma) \\ \rho_{(B-R)T_N,v_c} = 0.17^{+0.15}_{-0.15} & (p = 0.14; CL = 1.48 \sigma) \end{cases}, \quad (13)$$

indicating that none remains without the effect of the other variable, even if the case of $\rho_{(B-R)v_c,T_N}$ is rather close to 2.5σ .

To conclude, for CKBOs

- it is statistically equivalent to use ψ , ε , or v_c .
- the correlation between $(B - R)$ and perihelia may be simply an effect that arises because in the current sample of measured objects the perihelion is related with inclination, where inclination is the dominating variable. The same is true if we replace the variable inclination by the Tisserand parameter.
- the correlations between $(B - R)$ and both Tisserand parameter and inclination cannot be statistically separated from the close link between T_N and i . None dominates the other (see Fig. 4).
- although less clear than for the previous case, the correlation between $(B - R)$ and collisional velocity cannot be statistically separated from the dependence of v_c on inclination and on T_N .
- collisional velocity v_c , Tisserand parameter T_N , or orbital inclination i may be the true physically relevant variables linked with $(B - R)$ surface colors.
- given previous studies on the lack of evidence that surface colors are affected by v_c , it is most likely that only either T_N or i can be the variables that truly relate with colors —




Fig. 4. Top: $(B-R)$ vs. orbital inclination plot of CKBOs and all KBOs with indication of Haumea family objects. The correlation for CKBOs alone is stronger than for all KBOs. The clustering of Haumea family objects had a strong effect on this correlation. Middle: Tisserand parameter relative to Neptune (T_N) vs. orbital inclination as for the above. T_N is very strongly correlated with i for CKBOs. Bottom: $(B-R)$ vs. T_N as for the above. Given the very strong link between T_N and i , statistically one cannot know if only T_N or i have a (potentially) physically meaningful relation with $(B-R)$ but none dominates the other.

although causality cannot be assessed with this analysis —, and that dynamical history plays the central role.

It is interesting to see that when the peculiar Haumea family objects are removed from the analysis of CKBOs, the color-inclination correlation diminishes considerably (see Fig. 4). Peixinho et al. (2008) reported $\rho = -0.70$, for $n = 69$ objects, whereas now we have $\rho = -0.50$ for $n = 79$ objects, excluding the Haumea family. When Doressoundiram et al. (2005b) analyzed the correlation using two different classification schemes for CKBOs, one using $q > 35$ AU as cutoff and another using $q > 39$ AU, they reported $\rho = -0.65$ ($n = 27$) for the former and $\rho = -0.41$ ($n = 20$, and $SL < 2\sigma$) for the latter, four of the seven objects that had $q < 39$ AU are now known to be members of the Haumea family. In other words, without the Haumea family, the color-inclination correlation among CKBOs is no longer a strong correlation but a moderate one.

It is interesting to note that CKBOs do not show evidence for a correlation between $(B-V)$ vs. $(V-R)$, which is present in almost all the other more undersampled families. We return to this matter in Sect. 5.9.1.

5.7. All KBOs*

We have also analyzed the $n = 322$ KBOs* in our sample, that is all objects except Centaurs, Haumea family objects, and retrograde orbits. All color indexes correlate with each other, as expected from previous works. However, it is interesting to note that the color-inclination correlation for all KBOs, although weaker, is within the error-bars of the same correlation for CKBOs. This suggests that the colors of most non-CKBO families of KBOs are not totally uncorrelated with inclination, otherwise the correlation would be wiped out. The color-inclination correlation for $n = 186$ KBOs* removing the CKBOs is $\rho = -0.30^{+0.07}_{-0.07}$, $p = 0.000042$, $CL = 4.09\sigma$ (better than 3σ after FDR correction). We must keep in mind that if the population's ρ is -0.5 , to have only a 10% risk of failing to detect it at a 2.5σ level requires a sample of $n = 55$ objects, but if it is -0.3 , we need $n = 156$.

SDOs do have a sampling that should be enough to detect most correlations $\geq |\pm 0.5|$, but show $\rho_{(B-R)i} = -0.10^{+0.15}_{-0.15}$ ($n = 54$) with a p-value ≈ 0.50 . Plutinos have a smaller sampling and show a $\rho_{(B-R)i} = -0.30^{+0.17}_{-0.15}$ ($n = 45$), where the first correlation is below the 2.5σ level after correcting for FDR. For non-Plutino resonants we have $\rho_{(B-R)i} = -0.26^{+0.13}_{-0.12}$ ($n = 67$), but below a 2σ level after correcting for FDR. These results indicate that it is unlikely that any non-CKBO family will have a yet-to-be detected color-inclination correlation stronger than -0.5 , but all may have a correlation of ~ -0.3 for which their sampling is too low to warrant detection. Clearly, the $(B-R)$ vs. i trend is stronger among CKBOs, regardless of the current lower sampling of other families.

To verify this behavior, we have simulated 1000 samples with 79 (x, y) data-points correlated with $\rho = -0.50$ (corresponding to our CKBOs) and 189 (x, y) data-points randomly generated, that is, with $\rho \sim 0$ (corresponding to the rest of our KBOs). The simulations show a distribution with $\langle \rho \rangle_{-\sigma}^{+\sigma} = -0.17^{+0.05}_{-0.05}$, which is not compatible with the observed $\rho = -0.30^{+0.07}_{-0.07}$, hence the weak color-inclination correlation seen among KBOs* is not simply a result of its stronger presence among CKBOs.

Weak correlations of $\rho \sim \pm 0.2$ are detected between colors and perihelion and eccentricity. Although weak in magnitude and only accounting for $\sim 5\%$ of the color variability — ρ^2 may be roughly interpreted as the proportion of the variability of Y

that could be explained as a result of the variability of X — they are statistically significant, meaning that they are not a random effect.

5.8. Binary or multiple KBOs

Our sample has $n = 48$ binary or multiple objects, including Pluto and Haumea, and we have also analyzed the primaries of these systems as a group to see if they behave differently from the non-binaries.

Their color-color correlation does not show a different behavior from the previous cases. Their surface colors correlate both with T_N , with ν_c , and i . A total of 26 out of these 48 objects are CKBOs, which are known to correlate with the aforementioned variables. Only $n = 35$ binary or multiple KBOs possess measured $(B - R)$ colors, which means $n = 14$ CKBOs. Analyzing only the $n = 20$ non-CKBOs, also excluding Haumea, we obtain $\rho = -0.49^{+0.24}_{-0.18}$, $p = 0.027$, $CL = 2.21\sigma$ (without correcting for FDR). The result is not significant, and given the sample size, we have $\rho_{\beta 2.5\sigma} = -0.76$, which means that the subsampling is too small to confirm the existence of $\rho_{(B-R)i} \approx -0.50$, like for CKBOs, or $\rho_{(B-R)i} \approx -0.30$ like for non-CKBO KBOs as discussed above.

5.9. All* objects

As indicated in Sect. 2, we have removed from our global analysis objects belonging to the Haumea family and those with retrograde orbits. Our all* objects sample has $n = 354$ objects. The only relevant difference between the analysis of all* objects and that of KBOs* alone is the increase of the correlations between absolute magnitude H_R and both q and a . These two cases may be simply interpreted as a bias. That is, fainter objects, which are most often also smaller, are hard to observe at high semi-major axes or high perihelia. When we include Centaurs in the sample, which are mostly small and faint, this becomes even more obvious. Nonetheless, it is a known fact that there are no large Centaurs, and this is not a bias.

5.9.1. Consecutive color correlations

Throughout all publications on the correlation between colors of Centaurs and KBOs, we see the tendency that with increasing wavelength of the region covered by the colors in question, the correlations tend to diminish. For example, in all works the correlations between $(B - V)$ vs. $(V - R)$ have been shown to be always stronger in magnitude than the correlations between $(J - H)$ vs. $(H - K)$. However, it is also a fact that the error-bars of visible colors are smaller than those of near-IR colors. Might this be the reason for the observed degradation of the correlation for increasing wavelength or not?

To answer this question, we have analyzed the relative median error-bar of each color normalized to the 6σ range of the corresponding color. For example, the 6σ range of $(B - V)$ color for all* is 1.036, the median error-bar is 0.08, therefore, after normalization, if $(B - V)$ ranges from 0 to 1 the median error-bars would be 0.077. With this procedure we can use Eq. 4 and, with an assumption on the parent population ρ value, see the effect of the error-bars on the detectable ρ^* (see Sect. 3.3)

Since the correlation of $(B - V)$ vs. $(V - R)$ usually is $\sim 0.5 - 0.6$, we assume, for simplicity, two cases: i) the parent population ρ is 0.7 and it should be 0.7 regardless of the spectral region we examine, ii) the same as the previous, but with

Table 4. Spearman ρ for consecutive colors.

X vs. Y	n	Observed $\rho_{-\sigma}^{+\sigma}$	Error σ_o	Degraded $\rho_{0.7}^*$	Degraded $\rho_{0.6}^*$	Detectable $\rho_{\beta 2.5\sigma}$
All*						
$(B - V)(V - R)$	292	$0.54^{+0.06}_{-0.07}$	0.177	0.54	0.48	0.22
$(V - R)(R - I)$	280	$0.34^{+0.07}_{-0.08}$	0.192	0.52	0.47	0.23
KBOs*						
$(B - V)(V - R)$	266	$0.51^{+0.07}_{-0.07}$	0.179	0.53	0.48	0.23
$(V - R)(R - I)$	252	$0.30^{+0.08}_{-0.08}$	0.195	0.51	0.47	0.24
KBOs* except CKBOs						
$(B - V)(V - R)$	185	$0.61^{+0.07}_{-0.08}$	0.167	0.55	0.49	0.28
$(V - R)(R - I)$	175	$0.36^{+0.09}_{-0.10}$	0.179	0.53	0.48	0.29
Classical KBOs						
$(B - V)(V - R)$	78	$0.15^{+0.15}_{-0.16}$	0.234	0.47	0.43	0.43
$(V - R)(R - I)$	76	$0.07^{+0.16}_{-0.16}$	0.241	0.46	0.42	0.43
Other resonants						
$(B - V)(V - R)$	67	$0.68^{+0.09}_{-0.12}$	0.178	0.54	0.48	0.50
$(V - R)(R - I)$	68	$0.44^{+0.13}_{-0.15}$	0.182	0.53	0.48	0.56
SDOs						
$(B - V)(V - R)$	53	$0.60^{+0.11}_{-0.14}$	0.164	0.55	0.50	0.51
$(V - R)(R - I)$	45	$0.29^{+0.19}_{-0.22}$	0.175	0.54	0.49	0.56
Plutinos						
$(B - V)(V - R)$	45	$0.55^{+0.16}_{-0.21}$	0.186	0.53	0.47	0.56
$(V - R)(R - I)$	42	$0.30^{+0.18}_{-0.21}$	0.265	0.43	0.40	0.57
Binary or multiple KBOs						
$(B - V)(V - R)$	34	$0.55^{+0.15}_{-0.20}$	0.147	0.58	0.51	0.63
$(V - R)(R - I)$	35	$0.35^{+0.19}_{-0.23}$	0.139	0.59	0.52	0.65
Centaurs						
$(B - V)(V - R)$	26	$0.67^{+0.13}_{-0.18}$	0.119	0.61	0.54	0.71
$(V - R)(R - I)$	28	$0.61^{+0.15}_{-0.20}$	0.159	0.56	0.50	0.69
SDKBOs						
$(B - V)(V - R)$	20	$0.55^{+0.25}_{-0.40}$	0.171	0.55	0.49	0.70
$(V - R)(R - I)$	20	$0.16^{+0.31}_{-0.35}$	0.157	0.56	0.50	0.70

$\rho = 0.6$. Table 4 summarizes the results. For each X vs. Y pair of colors we have their own sampling n , observed ρ , and normalized median error-bars σ_o . From Eq. 4 we compute the degraded $\rho_{\text{Degraded}} \equiv \rho^*$ for both 0.7 and 0.6, and from the sampling n we solve Eq. 2 to obtain the detectable ρ at a 2.5σ level accepting only a risk $\beta = 0.1$ of missing it, $\rho_{\beta 2.5\sigma}$ (see Sect. 3.2). Statistically ρ decreases with higher error-bars which are also typically associated with higher wavelengths but the variation is small. Observationally it decreases at a much faster rate. Therefore, we may conclude that, globally, the $(B - V)$ vs. $(V - R)$ correlation is stronger than the $(V - R)$ vs. $(R - I)$ one, and it is not an effect created by larger error-bars, that is, the color-color correlations do diminish with increasing wavelength. This result does not discard the possibility that this cannot be the case for some specific subgroups of objects, however.

Examining these correlations for each family in more detail, we may see a similar behavior for all of them except for CKBOs. The sampling of Plutinos, binary or multiple KBOs, Centaurs, and SDKBOs does not warrant detection of correlations weaker than ~ 0.6 at a β risk of 10%.

Since it is clear that the $(V-R)$ vs. $(R-I)$ correlation is indeed weaker than that of $(B-V)$ vs. $(V-R)$, it is not surprising that we do not detect it among SDOs, Plutinos, binary or multiple KBOs, or SDKBOs. For the latter we do not even detect the $(B-V)$ vs. $(V-R)$ correlation. For Centaurs, both are detected and show similar strengths. As a result of the low sampling, however, the correlation error-bars are too high to conclude that they are indeed similar, in contrast to what we have seen in other objects.

The most interesting case, however, are CKBOs, since these objects show neither $(B-V)$ vs. $(V-R)$ nor $(V-R)$ vs. $(R-I)$ correlations (see Sect. 5.6.1). The visible spectrophotometric behavior of CKBOs, or at least one type of CKBOs, is distinct from other KBOs. We discuss it below.

5.9.2. Peculiar convex spectrophotometric behaviors

Surface spectra of Centaurs and KBOs tend to be straight in the visible and flatten toward the near-IR JHK bands, most particularly when the visible slope is red, although the near-IR behavior tends to be more erratic. Figures 2 and 3 show that in $(B-V)$ vs. $(V-R)$ the objects tend to lie in the vicinity of the reddening line, meaning that they possess rather straight spectra in the BVR wavelength range (pictorially: — or /), or above it, meaning that they possess a slightly convex spectra in that same range (pictorially: $\smile$). In the $(V-R)$ vs. $(R-I)$ plots we may see that objects are distributed above and below the reddening line. We have identified 11 objects that fall far below the BVR reddening line: 2000CL₁₀₄, 2002VT₁₃₀, 2001RZ₁₄₃, 1999JD₁₃₂, 2001HY₆₅, 2001HZ₅₈, 1997RT₅, 2000CL₁₀₅, 2002VD₁₃₁³ (all Classical KBOs), 2001KY₇₆ (Plutino), and 1999RJ₂₁₅ (SDO) — these are all mostly classical KBOs. Their normalized spectrophotometric reflectivity spectra are plotted in Fig. 5 along with the spectra resulting from the mean colors of all* objects (the error bars from the latter are the standard deviation of the color distribution and not the average error on each color). Spectra are normalized to 1 at the center of the V -band filter, and each one is shifted by 0.4 for clarity. The convex behavior these objects present in BVR might be a mere consequence of a concave (pictorially: $\frown$) behavior in VRI , that is, the existence of an absorption band at $\sim 6000-7000$ Å. It is tempting to associate this behavior with aqueous altered minerals creating absorption features in this region, as once seen with spectroscopy on (38628) Huya and (47932) 2000GN₁₇₁ by de Bergh et al. (2004), but the effect on visible colors is not expected to be very strong (see Barucci et al. 2008, for a review). The object 2001RZ₁₄₃ does seem to be fully convex up to the I band, and we do not possess $V-I$ colors of objects 2002VD₁₃₁, 2000CL₁₀₄, and 1999RJ₂₁₅ to see if the spectral slope changes its convexity. Nonetheless, with the exception of 2001RZ₁₄₃, the other 10 aforementioned objects seem good candidates to measure for the long-sought hydration features among Centaurs and KBOs or any other absorption feature in the visible spectrum.




Fig. 5. Spectrophotometric reflectivity of 11 peculiar objects exhibiting a convex behavior at $\sim 4500-6000$ Å. Spectra are normalized to 1 at $\lambda = 5454$ Å and are consecutively shifted by 0.4 for clarity. This apparent convexity may be created by a concave behavior at $\sim 6000-7000$ Å, which suggests some absorption feature on their surfaces in this wavelength range. The spectrum resulting from the mean $B-V$, $V-R$, and $V-I$ colors, and their standard deviations, for all* objects is shown in gray.

5.10. Bimodal colors of small objects

Peixinho et al. (2012) found that both Centaurs and KBOs present the so-called bimodal ($B-R$) visible color behavior at a p -value=0.001 (3.3σ) for absolute magnitudes $H_R(\alpha) \geq 6.8$. In a somewhat different manner, Fraser & Brown (2012) found that all small objects ($H_R(\alpha) \geq 6$) with perihelion $q < 35$ AU are bimodal, which will therefore include Centaurs. With this new dataset we have carried out the same analysis as Peixinho et al. (2012), obtaining a maximum of significance for the same bimodality for the $n = 143$ objects with $H_R(\alpha) \geq 6.82$, with a p -value=0.0046 (2.8σ). Although the evidence for a bimodal

³ Note that 2002VD₁₃₁ was not included in the statistical analysis of classical KBOs because of its extreme outlier behavior.




Fig. 6. $(B - R)$ colors vs. $H_R(\alpha)$ absolute magnitude (uncorrected for phase effects) for all objects in our database. Centaurs and KBOs are indicated with different symbols. Haumea family objects and the two retrograde objects are also highlighted. The N -shape reported by Peixinho et al. (2012) is clearly visible, and faint object bimodality is detected for $H_R(\alpha) \geq 6.82$, but at 2.8σ level (p-value=0.0046).

behavior is very strong, we do not find a 3σ result with the new dataset (see Fig. 6).

Testing the $n = 112$ objects with perihelion $q < 35$ AU and $H_R(\alpha) \geq 6.0$ from the all* sample results in a p-value=0.126 (1.5σ), which means that there is no evidence for bimodality.

Interestingly, although the bimodality among Centaurs seems clear to the eye, when testing only for the $n = 29$ Centaurs in the sample we obtain a p-value=0.018 (2.4σ).

6. Conclusions

We reanalyzed the correlations of $BVRI$ -band visible colors for $n = 354$ Centaurs and KBOs. The $(B - V)$, $(V - R)$, $(R - I)$, $(B - R)$, $(V - I)$, and $(B - I)$ color indexes were carefully computed from all the available measurements in the literature, discarding color indexes computed from two bands taken more than two hours apart, and rejecting clear outliers from the averaging process. The complete sample consisted of $n = 366$ objects, but we removed from the analysis $n = 2$ objects with retrograde orbits and $n = 10$ Haumea family objects. Nonetheless, these objects were also plotted in our figures.

- We used an algorithm for the Spearman-rank correlation — also known as Spearman- ρ — taking into account not only the data error-bars but also the error on the estimate due to the finite sampling and corrected the significance levels using the false-discovery rate (FDR) due to the large number of tests each sample was being subjected to.
- We analyzed for the first time the risk β of missing any possible correlations as a result of our limited sampling and the sampling necessary to ensure detection at a chosen significance level.
- With samples of $n < 70$ objects, assumed to be unbiased extractions of the parent population, we have a $\beta > 10\%$ risk of missing strong correlations, and if $n < 90$ medium strength (moderate) correlations may still be missed. When analyzing the traditional dynamical families, only classical

KBOs, as a whole, do not require larger sampling to detect presumably relevant correlations. If one aims to study the subfamilies of hot classical, cold classical, inner classicals, or other, then larger samples are still needed. Centaurs are particularly undersampled and should be more than doubled. Plutinos and SDOs would benefit from a 50% sample increase. Other resonants, as a whole, are not undersampled, but like the Plutinos, each resonant family should be analyzed as a single group, and each one of them is largely undersampled. The sampling of the objects we grouped as scattered or detached KBOs would also need to be tripled. Note that the detection requirements for sample-size vs. correlation strength are general and would need to be applied to any possible subfamily or distinct classification one decides to study. However, the samples are assumed to be unbiased. Including possible sample biasing effects would require a more complex analysis.

- We studied the degradation of a detectable correlation as a function of the data error-bars and concluded that after a certain correlation has been detected, it is pointless to increase the sampling while maintaining the same error-bars. Only by reducing the observational error-bars can a possibly stronger correlation be measured.
- We showed that the location of so-called reddening line in the color-color plots is quite sensitive to the central wavelengths of similar filters used on different telescopes. The effect is strong in the $(B - V)$ vs. $(V - R)$ plots, but weak in the $(V - R)$ vs. $(R - I)$ plots.
- The correlation between $(B - V)$ and $(V - R)$ colors is stronger than the correlation between $(V - R)$ and $(R - I)$, regardless of the observational error-bars.
- We analyzed the different parameters that have been used in the context of the color-inclination correlation of classical KBOs and the collisional resurfacing scenario using partial correlation tests. Statistically, it is equivalent to using the orbital excitation ϵ , Öpik's ψ , or the collisional velocity v_c . The detected color-perihelion correlation is most likely physically irrelevant and an effect of the strong relation between perihelion q and inclination i among CKBOs. Both orbital inclination i and Tisserand parameter relative to Neptune T_N are the most likely physically relevant parameters related with the color variation of CKBOs, therefore the relation is probably related to dynamical history alone. Statistically, the collisional velocity cannot be separated from its dependence on inclination i or T_N . Without the Haumea family, the color-inclination correlation of CKBOs is no longer strong, but moderate ($\rho \sim 0.5$).
- Although weaker than for CKBOs, a color-inclination correlation must still exist among the non-classical KBOs.
- Classical KBOs are the only family that show neither $(B - V)$ vs. $(V - R)$ nor $(V - R)$ vs. $(R - I)$ correlation.
- We identified ten objects, only two of which are not CKBOs, that exhibit a convex spectral behavior from their $(B - V)$ and $(V - R)$ colors and a concave behavior from their $(V - R)$ and $(R - I)$ colors, suggesting an absorption band or feature at $\sim 6000 - 7000 \text{ \AA}$, which renders them good candidates for detailed spectroscopic studies.
- We detected evidence for the $(B - R)$ color bimodality for objects with absolute magnitude $H_R(\alpha) \geq 6.82$ at a 2.8σ level (p-value=0.0046) — see Peixinho et al. (2012). Paradoxically, the evidence for bimodality among Centaurs reaches only 2.4σ — see Peixinho et al. (2003) —, and we find no evidence for bimodality of objects with perihelion $q < 35$ AU and $H_R(\alpha) \geq 6.0$ — see Fraser & Brown (2012).

We must stress that we analyzed the correlations and sampling requirements of the traditional dynamical families of KBOs and Centaurs as defined in Sect. 2, with the aim to characterize each of them, and with the underlying statistical assumption that each sample is or will be unbiased. The physical and/or chemical relevance of the dynamical classification system used was not discussed. Using distinct classification systems, subsystems, or groupings should lead to distinct correlations as well as to distinct sampling considerations, and such works are to be encouraged.

A similar study, extended to the near-IR colors, is under preparation.

Acknowledgements. We thank the anonymous referee for the comments that helped to improve this manuscript. We thank Audrey Thirouin for her LaTeX “magic”. NP acknowledges funding by the Gemini-Conicyt Fund, allocated to the project N° 32120036.

References

- Alvarez-Candal, A., Barucci, M. A., Merlin, F., et al. 2010, *A&A*, 511, A35
- Barucci, M. A., Belskaya, I. N., Fulchignoni, M., & Birlan, M. 2005, *AJ*, 130, 1291
- Barucci, M. A., Boehnhardt, H., Dotto, E., et al. 2002, *A&A*, 392, 335
- Barucci, M. A., Brown, M. E., Emery, J. P., & Merlin, F. 2008, *Composition and Surface Properties of Transneptunian Objects and Centaurs*, ed. M. A. Barucci, H. Boehnhardt, D. P. Cruikshank, A. Morbidelli, & R. Dotson, 143–160
- Barucci, M. A., Doressoundiram, A., Tholen, D., Fulchignoni, M., & Lazzarin, M. 1999, *Icarus*, 142, 476
- Barucci, M. A., Romon, J., Doressoundiram, A., & Tholen, D. J. 2000, *Astron. J.*, 120, 496
- Batygin, K., Brown, M. E., & Fraser, W. C. 2011, *ApJ*, 738, 13
- Bauer, J. M., Meech, K. J., Fernández, Y. R., et al. 2003, *Icarus*, 166, 195
- Benecchi, S. D., Noll, K. S., Grundy, W. M., et al. 2009, *Icarus*, 200, 292
- Benecchi, S. D., Noll, K. S., Stephens, D. C., Grundy, W. M., & Rawlins, J. 2011, *Icarus*, 213, 693
- Benjamini, Y. & Hochberg, Y. 1995, *Journal of the Royal Statistical Society. Series B (Methodological)*, 57, 289
- Boehnhardt, H., Bagnulo, S., Muinonen, K., et al. 2004, *A&A*, 415, L21
- Boehnhardt, H., Delsanti, A., Barucci, A., et al. 2002, *A&A*, 395, 297
- Boehnhardt, H., Tozzi, G. P., Birkle, K., et al. 2001, *A&A*, 378, 653
- Bonett, D. G. & Wright, T. A. 2000, *Psychometrika*, 65, 23
- Bonferroni, C. E. 1935, *Il calcolo delle assicurazioni su gruppi di teste*, ed. Rome, Italy, 13–60
- Brown, M. E., Barkume, K. M., Ragozzine, D., & Schaller, E. L. 2007, *Nature*, 446, 294
- Brown, M. E., Schaller, E. L., & Fraser, W. C. 2011, *ApJ*, 739, L60
- Dalle Ore, C. M., Dalle Ore, L. V., Roush, T. L., et al. 2013, *Icarus*, 222, 307
- de Bergh, C., Boehnhardt, H., Barucci, M. A., et al. 2004, *A&A*, 416, 791
- de Bergh, C., Delsanti, A., Tozzi, G. P., et al. 2005, *A&A*, 437, 1115
- Delsanti, A., Hainaut, O., Jourdeuil, E., et al. 2004, *A&A*, 417, 1145
- Delsanti, A., Peixinho, N., Boehnhardt, H., et al. 2006, *AJ*, 131, 1851
- Delsanti, A. C., Boehnhardt, H., Barrera, L., et al. 2001, *A&A*, 380, 347
- DeMeo, F. E., Barucci, M. A., Merlin, F., et al. 2010, *A&A*, 521, A35
- DeMeo, F. E., Fornasier, S., Barucci, M. A., et al. 2009, *A&A*, 493, 283
- Doressoundiram, A., Barucci, M. A., Romon, J., & Veillet, C. 2001, *Icarus*, 154, 277
- Doressoundiram, A., Barucci, M. A., Tozzi, G. P., et al. 2005a, *Planet. Space Sci.*, 53, 1501
- Doressoundiram, A., Boehnhardt, H., Tegler, S. C., & Trujillo, C. 2008, *Color Properties and Trends of the Transneptunian Objects*, ed. M. A. Barucci, H. Boehnhardt, D. P. Cruikshank, A. Morbidelli, & R. Dotson, 91–104
- Doressoundiram, A., Peixinho, N., de Bergh, C., et al. 2002, *AJ*, 124, 2279
- Doressoundiram, A., Peixinho, N., Doucet, C., et al. 2005b, *Icarus*, 174, 90
- Doressoundiram, A., Peixinho, N., Moullet, A., et al. 2007, *AJ*, 134, 2186
- Dotto, E., Barucci, M. A., Boehnhardt, H., et al. 2003, *Icarus*, 162, 408
- Edgeworth, K. E. 1943, *Journal of the British Astronomical Association*, 53, 181
- Edgeworth, K. E. 1949, *Mon. Not. R. Astron. Soc.*, 109, 600
- Efron, B. & Tibshirani, R. J. 1993, *An Introduction to the Bootstrap* (Chapman & Hall/CRC)
- Elliot, J. L., Kern, S. D., Clancy, K. B., et al. 2005, *AJ*, 129, 1117
- Emel’yanenko, V. V., Asher, D. J., & Bailey, M. E. 2013, *Earth Moon and Planets*, 110, 105
- Fernandez, J. A. 1980, *MNRAS*, 192, 481
- Ferrin, I., Rabinowitz, D., Schaefer, B., et al. 2001, *ApJ*, 548, L243
- Fisher, R. A. 1915, *Biometrika*, 10, pp. 507
- Fornasier, S., Doressoundiram, A., Tozzi, G. P., et al. 2004, *A&A*, 421, 353
- Fouchard, M., Rickman, H., Froeschlé, C., & Valsecchi, G. B. 2014, *Icarus*, 231, 99
- Fraser, W. C. & Brown, M. E. 2012, *ApJ*, 749, 33
- Gil-Hutton, R. 2002, *Planet. Space Sci.*, 50, 57
- Gil-Hutton, R. & Licandro, J. 2001, *Icarus*, 152, 246
- Gladman, B., Marsden, B. G., & Vanlaerhoven, C. 2008, *Nomenclature in the Outer Solar System*, ed. Barucci, M. A., Boehnhardt, H., Cruikshank, D. P., Morbidelli, A., & Dotson, R., 43–57
- Gomes, R. S. 2003, *Icarus*, 161, 404
- Green, S. F., McBride, N., O’Ceallaigh, D. P., et al. 1997, *MNRAS*, 290, 186
- Gulbis, A. A. S., Elliot, J. L., & Kane, J. F. 2006, *Icarus*, 183, 168
- Hainaut, O. R., Boehnhardt, H., & Protapapa, S. 2012, *A&A*, 546, A115
- Hainaut, O. R., Delahodde, C. E., Boehnhardt, H., et al. 2000, *A&A*, 356, 1076
- Hainaut, O. R. & Delsanti, A. C. 2002, *A&A*, 389, 641
- Horner, J., Evans, N. W., & Bailey, M. E. 2004, *MNRAS*, 354, 798
- Jewitt, D. & Luu, J. 1993, *Nature*, 362, 730
- Jewitt, D. & Luu, J. 1998, *AJ*, 115, 1667
- Jewitt, D., Peixinho, N., & Hsieh, H. H. 2007, *AJ*, 134, 2046
- Jewitt, D. C. & Luu, J. X. 2001, *AJ*, 122, 2099
- Kaňuchová, Z., Brunetto, R., Melita, M., & Strazzulla, G. 2012, *Icarus*, 221, 12
- Kavelaars, J., Jones, L., Gladman, B., Parker, J. W., & Petit, J.-M. 2008, *The Orbital and Spatial Distribution of the Kuiper Belt*, ed. M. A. Barucci, H. Boehnhardt, D. P. Cruikshank, A. Morbidelli, & R. Dotson, 59–69
- Kendall, M. G., Kendall, S. F. H., & Smith, B. B. 1939, *Biometrika*, 30, pp. 251
- Kresák, L. 1972, in *IAU Symposium*, Vol. 45, *The Motion, Evolution of Orbits, and Origin of Comets*, ed. G. A. Chebotarev, E. I. Kazimirschak-Polonskaia, & B. G. Marsden, 503
- Kuiper, G. P. 1951, in *Proceedings of a topical symposium, commemorating the 50th anniversary of the Yerkes Observatory and half a century of progress in astrophysics*, ed. J. Hynek (New York: McGraw-Hill), 357
- Lacerda, P., Fornasier, S., Lellouch, E., et al. 2014, *ApJ*, 793, L2
- Lazzaro, D., Florczak, M. A., Angeli, C. A., et al. 1997, *Planet. Space Sci.*, 45, 1607
- Leonard, F. C. 1930, *Leaflet of the Astronomical Society of the Pacific*, 1, 121
- Levenberg, K. 1944, *Quarterly Journal of Applied Mathematics*, II, 164
- Levison, H. F. 1996, in *ASP Conf. Ser. 107: Completing the Inventory of the Solar System*, 173–191
- Levison, H. F. & Duncan, M. J. 1997, *Icarus*, 127, 13
- Levison, H. F., Morbidelli, A., Tsiganis, K., Nesvorný, D., & Gomes, R. 2011, *AJ*, 142, 152
- Luu, J. & Jewitt, D. 1996, *AJ*, 112, 2310
- Lykawka, P. S. & Mukai, T. 2007, *Icarus*, 189, 213
- Macklin, J. T. 1982, *Mon. Not. R. Astron. Soc.*, 199, 1119
- Marquardt, D. W. 1963, *Journal of the Society for Industrial and Applied Mathematics*, 11, 431
- Miller, R. G. 1981, *Simultaneous Statistical Inference* (Springer New York, 2nd ed.)
- Modarres, R. 1996, *Statistics and Computing*, 6, 139
- Öpik, E. J. 1976, *Interplanetary encounters: close-range gravitational interactions*. (Elsevier Science Ltd)
- Peixinho, N., Boehnhardt, H., Belskaya, I., et al. 2004, *Icarus*, 170, 153
- Peixinho, N., Delsanti, A., Guilbert-Lepoutre, A., Gafeira, R., & Lacerda, P. 2012, *A&A*, 546, A86
- Peixinho, N., Doressoundiram, A., Delsanti, A., et al. 2003, *A&A*, 410, L29
- Peixinho, N., Lacerda, P., & Jewitt, D. 2008, *AJ*, 136, 1837
- Peixinho, N., Lacerda, P., Ortiz, J. L., et al. 2001, *A&A*, 371, 753
- Perna, D., Barucci, M. A., Fornasier, S., et al. 2010, *A&A*, 510, A53
- Petit, J.-M., Kavelaars, J. J., Gladman, B. J., et al. 2011, *AJ*, 142, 131
- Rabinowitz, D. L., Barkume, K., Brown, M. E., et al. 2006, *ApJ*, 639, 1238
- Rabinowitz, D. L., Schaefer, B. E., Schaefer, M., & Tourtellotte, S. W. 2008, *AJ*, 136, 1502
- Rabinowitz, D. L., Schaefer, B. E., & Tourtellotte, S. W. 2007, *AJ*, 133, 26
- Ramírez, I., Michel, R., Sefako, R., et al. 2012, *ApJ*, 752, 5
- Romanishin, W., Tegler, S. C., & Consolmagno, G. J. 2010, *AJ*, 140, 29
- Romanishin, W., Tegler, S. C., Levine, J., & Butler, N. 1997, *AJ*, 113, 1893
- Romon-Martin, J., Barucci, M. A., de Bergh, C., et al. 2002, *Icarus*, 160, 59
- Romon-Martin, J., Delahodde, C., Barucci, M. A., de Bergh, C., & Peixinho, N. 2003, *A&A*, 400, 369
- Santos-Sanz, P., Lellouch, E., Fornasier, S., et al. 2012, *A&A*, 541, A92
- Santos-Sanz, P., Ortiz, J. L., Barrera, L., & Boehnhardt, H. 2009, *A&A*, 494, 693
- Sheppard, S. S. 2010, *AJ*, 139, 1394
- Sheppard, S. S. 2012, *AJ*, 144, 169
- Snodgrass, C., Carry, B., Dumas, C., & Hainaut, O. 2010, *A&A*, 511, A72
- Spearman, C. 1904, *Am. J. Psychol.*, 57, 72
- Stern, S. A. 2002, *AJ*, 124, 2297
- Tegler, S. C. & Romanishin, W. 1997, *Icarus*, 126, 212
- Tegler, S. C. & Romanishin, W. 1998, *Nature*, 392, 49
- Tegler, S. C. & Romanishin, W. 2000, *Nature*, 407, 979
- Tegler, S. C. & Romanishin, W. 2003, *Icarus*, 161, 181
- Tegler, S. C., Romanishin, W., & Consolmagno, G. J. 2003, *ApJ*, 599, L49
- Thébault, P. & Doressoundiram, A. 2003, *Icarus*, 162, 27
- Thébault, P. 2003, *Earth Moon and Planets*, 92, 233
- Tisserand, F. 1896, *Traité de Mécanique Céleste* (Tome IV) (Gauthier-Villars, Paris)
- Trujillo, C. A. & Brown, M. E. 2002, *ApJ*, 566, L125

Table 2. Significant Spearman ρ for each family and group of families

FDR correction ^a											
X^b	vs.	Y^c	n^d	$\rho_{-\sigma}^{+\sigma e}$	P-value ^f	$CL(\sigma)^g$	i^h	$\frac{i}{m} 2.5 \sigma^i$	$\frac{i}{m} 3 \sigma^j$	$\rho_{\beta 2.5\sigma}^k$	$\rho_{\beta 3\sigma}^l$
Centaurs											
$(B - V)$	vs.	$(B - R)$	26	$0.83^{+0.08}_{-0.13}$	$< 10^{-6}$	5.42	1	0.000135	0.000029	0.69	0.75
<i>Grt</i>	vs.	$(B - R)$	28	$0.82^{+0.08}_{-0.13}$	$< 10^{-6}$	5.42	2	0.000270	0.000059	0.67	0.73
<i>Grt</i>	vs.	$(V - I)$	28	$0.81^{+0.10}_{-0.20}$	$< 10^{-6}$	5.22	3	0.000404	0.000088	0.67	0.73
<i>Grt</i>	vs.	$(V - R)$	31	$0.81^{+0.09}_{-0.15}$	$< 10^{-6}$	> 8.0	4	0.000539	0.000117	0.64	0.71
$(V - R)$	vs.	$(B - R)$	28	$0.76^{+0.10}_{-0.15}$	0.000003	4.69	5	0.000674	0.000147	0.67	0.73
$(B - I)$	vs.	$(B - R)$	19	$0.84^{+0.09}_{-0.17}$	0.000007	4.49	6	0.000809	0.000176	0.77	0.83
$(V - I)$	vs.	$(B - R)$	25	$0.76^{+0.11}_{-0.19}$	0.000009	4.44	7	0.000943	0.000205	0.70	0.76
$(V - R)$	vs.	$(V - I)$	28	$0.73^{+0.12}_{-0.20}$	0.000011	4.39	8	0.001078	0.000235	0.67	0.73
<i>Grt</i>	vs.	$(R - I)$	28	$0.72^{+0.12}_{-0.18}$	0.000015	4.33	9	0.001213	0.000264	0.67	0.73
<i>Grt</i>	vs.	$(B - V)$	26	$0.74^{+0.10}_{-0.15}$	0.000016	4.31	10	0.001348	0.000293	0.69	0.75
$(V - I)$	vs.	$(B - I)$	19	$0.82^{+0.10}_{-0.21}$	0.000016	4.32	11	0.001483	0.000323	0.77	0.83
$(R - I)$	vs.	$(V - I)$	28	$0.70^{+0.14}_{-0.23}$	0.000037	4.12	12	0.001617	0.000352	0.67	0.73
$(B - V)$	vs.	$(B - I)$	19	$0.80^{+0.11}_{-0.20}$	0.000041	4.10	13	0.001752	0.000382	0.77	0.83
<i>Grt</i>	vs.	$(B - I)$	19	$0.79^{+0.12}_{-0.23}$	0.000059	4.02	14	0.001887	0.000411	0.77	0.83
$(R - I)$	vs.	$(B - R)$	25	$0.69^{+0.13}_{-0.20}$	0.000118	3.85	15	0.002022	0.000440	0.70	0.76
$(B - V)$	vs.	$(V - R)$	26	$0.67^{+0.13}_{-0.18}$	0.000208	3.71	16	0.002157	0.000470	0.69	0.75
$(B - V)$	vs.	$(V - I)$	23	$0.69^{+0.13}_{-0.20}$	0.000237	3.68	17	0.002291	0.000499	0.72	0.78
— 3 σ FDR correction cutoff —											
$(V - R)$	vs.	$(R - I)$	28	$0.61^{+0.15}_{-0.20}$	0.000569	3.45	18	0.002426	0.000528	0.67	0.73
$(B - V)$	vs.	$(R - I)$	23	$0.66^{+0.15}_{-0.22}$	0.000582	3.44	19	0.002561	0.000558	0.72	0.78
$(R - I)$	vs.	$(B - I)$	19	$0.71^{+0.16}_{-0.30}$	0.000616	3.42	20	0.002696	0.000587	0.77	0.83
$(V - R)$	vs.	$(B - I)$	19	$0.71^{+0.14}_{-0.23}$	0.000664	3.40	21	0.002830	0.000616	0.77	0.83
— 2.5 σ FDR correction cutoff —											
Scattered disk objects											
$(B - I)$	vs.	$(B - R)$	40	$0.83^{+0.09}_{-0.18}$	$< 10^{-6}$	> 8.0	1	0.000148	0.000032	0.58	0.64
$(B - V)$	vs.	$(B - I)$	42	$0.75^{+0.10}_{-0.16}$	$< 10^{-6}$	> 8.0	2	0.000295	0.000064	0.56	0.63
$(B - V)$	vs.	$(B - R)$	51	$0.79^{+0.07}_{-0.10}$	$< 10^{-6}$	> 8.0	3	0.000443	0.000096	0.52	0.58
<i>Grt</i>	vs.	$(B - I)$	42	$0.80^{+0.07}_{-0.11}$	$< 10^{-6}$	> 8.0	4	0.000590	0.000129	0.56	0.63
<i>Grt</i>	vs.	$(B - R)$	51	$0.77^{+0.07}_{-0.10}$	$< 10^{-6}$	> 8.0	5	0.000738	0.000161	0.52	0.58
<i>Grt</i>	vs.	$(B - V)$	53	$0.64^{+0.10}_{-0.13}$	$< 10^{-6}$	5.12	6	0.000886	0.000193	0.51	0.57
<i>Grt</i>	vs.	$(V - I)$	48	$0.70^{+0.12}_{-0.18}$	$< 10^{-6}$	> 8.0	7	0.001033	0.000225	0.53	0.59
<i>Grt</i>	vs.	$(V - R)$	56	$0.73^{+0.11}_{-0.16}$	$< 10^{-6}$	> 8.0	8	0.001181	0.000257	0.50	0.55
$(V - I)$	vs.	$(B - I)$	42	$0.74^{+0.10}_{-0.14}$	$< 10^{-6}$	> 8.0	9	0.001329	0.000289	0.56	0.63
$(V - R)$	vs.	$(B - I)$	42	$0.73^{+0.11}_{-0.16}$	$< 10^{-6}$	> 8.0	10	0.001476	0.000321	0.56	0.63
$(V - R)$	vs.	$(B - R)$	51	$0.75^{+0.08}_{-0.10}$	$< 10^{-6}$	> 8.0	11	0.001624	0.000354	0.52	0.58
$(V - I)$	vs.	$(B - R)$	43	$0.67^{+0.12}_{-0.17}$	0.000001	4.97	12	0.001771	0.000386	0.56	0.62
$(B - V)$	vs.	$(V - R)$	53	$0.60^{+0.11}_{-0.14}$	0.000002	4.77	13	0.001919	0.000418	0.51	0.57
$(V - R)$	vs.	$(V - I)$	48	$0.60^{+0.14}_{-0.20}$	0.000007	4.49	14	0.002067	0.000450	0.53	0.59
$(B - V)$	vs.	$(V - I)$	45	$0.60^{+0.13}_{-0.18}$	0.000015	4.32	15	0.002214	0.000482	0.55	0.61
H_R	vs.	q	55	$-0.50^{+0.13}_{-0.11}$	0.000114	3.86	16	0.002362	0.000514	0.50	0.56
$(R - I)$	vs.	$(V - I)$	45	$0.52^{+0.16}_{-0.21}$	0.000284	3.63	17	0.002510	0.000546	0.55	0.61
$(R - I)$	vs.	$(B - I)$	40	$0.54^{+0.16}_{-0.22}$	0.000298	3.62	18	0.002657	0.000579	0.58	0.64
— 3 σ FDR correction cutoff —											
<i>Grt</i>	vs.	$(R - I)$	45	$0.46^{+0.16}_{-0.19}$	0.001317	3.21	19	0.002805	0.000611	0.55	0.61
— 2.5 σ FDR correction cutoff —											
Scattered or detached objects											
<i>Grt</i>	vs.	$(B - I)$	18	$0.82^{+0.12}_{-0.30}$	0.000033	4.15	1	0.000163	0.000036	0.79	0.84
— 3 σ FDR correction cutoff —											
$(B - I)$	vs.	$(B - R)$	18	$0.79^{+0.12}_{-0.25}$	0.000085	3.93	2	0.000326	0.000071	0.79	0.84
<i>Grt</i>	vs.	$(V - I)$	21	$0.73^{+0.14}_{-0.26}$	0.000149	3.79	3	0.000489	0.000107	0.75	0.81
<i>Grt</i>	vs.	$(B - R)$	20	$0.75^{+0.12}_{-0.20}$	0.000163	3.77	4	0.000653	0.000142	0.76	0.82
<i>Grt</i>	vs.	$(V - R)$	21	$0.72^{+0.15}_{-0.28}$	0.000223	3.69	5	0.000816	0.000178	0.75	0.81
continued on next page											

continued on next page

Table 2. continued.

X^b	vs.	Y^c	n^d	$\rho_{-\sigma}^{+\sigma e}$	P-value ^f	$CL(\sigma)^g$	i^h	FDR correction ^a			
								$\frac{i}{m} 2.5 \sigma^i$	$\frac{i}{m} 3 \sigma^j$	$\rho_{\beta 2.5\sigma}^k$	$\rho_{\beta 3\sigma}^l$
(V – R)	vs.	(B – R)	20	$0.73^{+0.16}_{-0.32}$	0.000257	3.65	6	0.000979	0.000213	0.76	0.82
(V – I)	vs.	(B – I)	18	$0.75^{+0.14}_{-0.26}$	0.000352	3.57	7	0.001142	0.000249	0.79	0.84
(B – V)	vs.	(B – R)	20	$0.71^{+0.17}_{-0.36}$	0.000396	3.54	8	0.001305	0.000284	0.76	0.82
(V – R)	vs.	(B – I)	18	$0.71^{+0.18}_{-0.36}$	0.001018	3.29	9	0.001468	0.000320	0.79	0.84
— 2.5 σ FDR correction cutoff —											
Plutinos											
Grt	vs.	(B – I)	18	$0.82^{+0.12}_{-0.30}$	0.000033	4.15	1	0.000163	0.000036	0.79	0.84
(B – I)	vs.	(B – R)	18	$0.79^{+0.12}_{-0.25}$	0.000085	3.93	2	0.000326	0.000071	0.79	0.84
Grt	vs.	(V – I)	21	$0.73^{+0.14}_{-0.26}$	0.000149	3.79	3	0.000489	0.000107	0.75	0.81
Grt	vs.	(B – R)	20	$0.75^{+0.12}_{-0.20}$	0.000163	3.77	4	0.000653	0.000142	0.76	0.82
Grt	vs.	(V – R)	21	$0.72^{+0.15}_{-0.28}$	0.000223	3.69	5	0.000816	0.000178	0.75	0.81
(V – R)	vs.	(B – R)	20	$0.73^{+0.16}_{-0.32}$	0.000257	3.65	6	0.000979	0.000213	0.76	0.82
(V – I)	vs.	(B – I)	18	$0.75^{+0.14}_{-0.26}$	0.000352	3.57	7	0.001142	0.000249	0.79	0.84
(B – V)	vs.	(B – R)	20	$0.71^{+0.17}_{-0.36}$	0.000396	3.54	8	0.001305	0.000284	0.76	0.82
(V – R)	vs.	(B – I)	18	$0.71^{+0.18}_{-0.36}$	0.001018	3.29	9	0.001468	0.000320	0.79	0.84
(B – V)	vs.	(B – I)	18	$0.67^{+0.17}_{-0.30}$	0.002264	3.05	10	0.001632	0.000355	0.79	0.84
(V – I)	vs.	(B – R)	20	$0.62^{+0.18}_{-0.27}$	0.003723	2.90	11	0.001795	0.000391	0.76	0.82
(V – R)	vs.	(V – I)	21	$0.58^{+0.19}_{-0.28}$	0.006273	2.73	12	0.001958	0.000426	0.75	0.81
(R – I)	vs.	(V – I)	21	$0.57^{+0.18}_{-0.26}$	0.006444	2.72	13	0.002121	0.000462	0.75	0.81
Grt	vs.	(B – V)	20	$0.56^{+0.18}_{-0.25}$	0.009851	2.58	14	0.002284	0.000497	0.76	0.82
(B – V)	vs.	(V – R)	20	$0.55^{+0.25}_{-0.40}$	0.011730	2.52	15	0.002447	0.000533	0.76	0.82
(B – R)	vs.	v_c	20	$-0.54^{+0.26}_{-0.18}$	0.014885	2.44	16	0.002611	0.000568	0.76	0.82
(B – V)	vs.	(V – I)	20	$0.51^{+0.20}_{-0.28}$	0.023125	2.27	17	0.002774	0.000604	0.76	0.82
— 3 σ FDR correction cutoff —											
(V – I)	vs.	v_c	22	$-0.45^{+0.28}_{-0.21}$	0.035165	2.11	18	0.002937	0.000639	0.74	0.80
(R – I)	vs.	(B – I)	18	$0.50^{+0.23}_{-0.33}$	0.035576	2.10	19	0.003100	0.000675	0.79	0.84
(V – I)	vs.	i	22	$-0.45^{+0.28}_{-0.21}$	0.035680	2.10	20	0.003263	0.000711	0.74	0.80
— 2.5 σ FDR correction cutoff —											
Other resonants											
B – I	vs.	B – R	62	$0.86^{+0.06}_{-0.11}$	$< 10^{-6}$	> 8.0	1	0.000163	0.000036	0.47	0.53
(B – V)	vs.	(B – I)	62	$0.81^{+0.06}_{-0.08}$	$< 10^{-6}$	> 8.0	2	0.000326	0.000071	0.47	0.53
(B – V)	vs.	(B – R)	67	$0.80^{+0.09}_{-0.14}$	$< 10^{-6}$	> 8.0	3	0.000489	0.000107	0.45	0.51
(B – V)	vs.	(V – I)	64	$0.69^{+0.08}_{-0.10}$	$< 10^{-6}$	> 8.0	4	0.000653	0.000142	0.46	0.52
(B – V)	vs.	(V – R)	67	$0.68^{+0.09}_{-0.12}$	$< 10^{-6}$	> 8.0	5	0.000816	0.000178	0.45	0.51
Grt	vs.	(B – I)	62	$0.86^{+0.06}_{-0.09}$	$< 10^{-6}$	> 8.0	6	0.000979	0.000213	0.47	0.53
Grt	vs.	(B – R)	67	$0.83^{+0.06}_{-0.09}$	$< 10^{-6}$	> 8.0	7	0.001142	0.000249	0.45	0.51
Grt	vs.	(B – V)	67	$0.75^{+0.07}_{-0.09}$	$< 10^{-6}$	> 8.0	8	0.001305	0.000284	0.45	0.51
Grt	vs.	(R – I)	68	$0.61^{+0.11}_{-0.14}$	$< 10^{-6}$	> 8.0	9	0.001468	0.000320	0.45	0.51
Grt	vs.	(V – I)	68	$0.79^{+0.07}_{-0.10}$	$< 10^{-6}$	> 8.0	10	0.001632	0.000355	0.45	0.51
Grt	vs.	(V – R)	71	$0.75^{+0.08}_{-0.11}$	$< 10^{-6}$	> 8.0	11	0.001795	0.000391	0.44	0.50
(R – I)	vs.	(B – I)	62	$0.70^{+0.08}_{-0.10}$	$< 10^{-6}$	> 8.0	12	0.001958	0.000426	0.47	0.53
(R – I)	vs.	(V – I)	68	$0.65^{+0.10}_{-0.13}$	$< 10^{-6}$	> 8.0	13	0.002121	0.000462	0.45	0.51
(V – I)	vs.	(B – I)	62	$0.83^{+0.06}_{-0.08}$	$< 10^{-6}$	> 8.0	14	0.002284	0.000497	0.47	0.53
(V – I)	vs.	(B – R)	64	$0.75^{+0.08}_{-0.10}$	$< 10^{-6}$	> 8.0	15	0.002447	0.000533	0.46	0.52
(V – R)	vs.	(B – I)	62	$0.75^{+0.08}_{-0.11}$	$< 10^{-6}$	> 8.0	16	0.002611	0.000568	0.47	0.53
(V – R)	vs.	(B – R)	67	$0.76^{+0.10}_{-0.15}$	$< 10^{-6}$	> 8.0	17	0.002774	0.000604	0.45	0.51
(V – R)	vs.	(V – I)	68	$0.66^{+0.09}_{-0.12}$	$< 10^{-6}$	> 8.0	18	0.002937	0.000639	0.45	0.51
(R – I)	vs.	(B – R)	64	$0.55^{+0.10}_{-0.12}$	0.000003	4.70	19	0.003100	0.000675	0.46	0.52
(B – V)	vs.	(R – I)	64	$0.53^{+0.10}_{-0.12}$	0.000006	4.53	20	0.003263	0.000711	0.46	0.52
(V – R)	vs.	(R – I)	68	$0.44^{+0.13}_{-0.15}$	0.000197	3.72	21	0.003426	0.000746	0.45	0.51
— 3 σ FDR correction cutoff —											
H_R	vs.	q	71	$-0.42^{+0.10}_{-0.09}$	0.000246	3.67	22	0.003589	0.000782	0.44	0.50
— 2.5 σ FDR correction cutoff —											
Classical KBOs											

continued on next page

Table 2. continued.

X^b	vs.	Y^c	n^d	$\rho_{-\sigma}^{+\sigma e}$	P-value ^f	$CL(\sigma)^g$	i^h	FDR correction ^a			
								$\frac{i}{m} 2.5 \sigma^i$	$\frac{i}{m} 3 \sigma^j$	$\rho_{\beta 2.5\sigma}^k$	$\rho_{\beta 3\sigma}^l$
(B – R)	vs.	ε	79	$-0.53^{+0.12}_{-0.10}$	$< 10^{-6}$	5.06	1	0.000124	0.000027	0.42	0.47
(B – R)	vs.	ψ	79	$-0.54^{+0.12}_{-0.10}$	$< 10^{-6}$	5.12	2	0.000248	0.000054	0.42	0.47
(B – R)	vs.	ν_c	79	$-0.54^{+0.11}_{-0.10}$	$< 10^{-6}$	5.12	3	0.000372	0.000081	0.42	0.47
(B – V)	vs.	(B – R)	78	$0.55^{+0.12}_{-0.10}$	$< 10^{-6}$	5.22	4	0.000496	0.000108	0.42	0.47
Grt	vs.	(V – R)	94	$0.54^{+0.15}_{-0.12}$	$< 10^{-6}$	> 8.0	5	0.000620	0.000135	0.39	0.43
(B – I)	vs.	(B – R)	53	$0.60^{+0.11}_{-0.15}$	0.000002	4.78	6	0.000744	0.000162	0.51	0.57
(B – R)	vs.	i	79	$-0.50^{+0.13}_{-0.11}$	0.000002	4.73	7	0.000868	0.000189	0.42	0.47
(B – R)	vs.	T_N	79	$0.50^{+0.13}_{-0.11}$	0.000003	4.71	8	0.000992	0.000216	0.42	0.47
Grt	vs.	(V – I)	80	$0.48^{+0.12}_{-0.13}$	0.000006	4.53	9	0.001116	0.000243	0.42	0.47
Grt	vs.	(B – R)	78	$0.47^{+0.14}_{-0.12}$	0.000017	4.30	10	0.001240	0.000270	0.42	0.47
(B – V)	vs.	ε	78	$-0.45^{+0.12}_{-0.10}$	0.000041	4.10	11	0.001364	0.000297	0.42	0.47
(B – V)	vs.	ν_c	78	$-0.44^{+0.12}_{-0.10}$	0.000052	4.05	12	0.001488	0.000324	0.42	0.47
(B – V)	vs.	ψ	78	$-0.44^{+0.12}_{-0.10}$	0.000063	4.00	13	0.001612	0.000351	0.42	0.47
(V – R)	vs.	T_N	97	$0.39^{+0.11}_{-0.12}$	0.000084	3.93	14	0.001736	0.000378	0.38	0.43
(B – V)	vs.	(B – I)	53	$0.51^{+0.15}_{-0.18}$	0.000104	3.88	15	0.001860	0.000405	0.51	0.57
Grt	vs.	ν_c	94	$-0.38^{+0.11}_{-0.10}$	0.000159	3.78	16	0.001984	0.000432	0.39	0.43
Grt	vs.	T_N	94	$0.38^{+0.10}_{-0.11}$	0.000167	3.76	17	0.002108	0.000459	0.39	0.43
Grt	vs.	ψ	94	$-0.38^{+0.11}_{-0.10}$	0.000168	3.76	18	0.002232	0.000486	0.39	0.43
(V – R)	vs.	(B – R)	78	$0.41^{+0.14}_{-0.16}$	0.000186	3.74	19	0.002356	0.000513	0.42	0.47
Grt	vs.	ε	94	$-0.38^{+0.11}_{-0.10}$	0.000187	3.74	20	0.002480	0.000540	0.39	0.43
Grt	vs.	(B – I)	53	$0.49^{+0.14}_{-0.17}$	0.000205	3.71	21	0.002604	0.000567	0.51	0.57
(V – R)	vs.	ψ	97	$-0.36^{+0.12}_{-0.11}$	0.000264	3.65	22	0.002728	0.000594	0.38	0.43
(B – V)	vs.	i	78	$-0.40^{+0.13}_{-0.12}$	0.000265	3.65	23	0.002852	0.000621	0.42	0.47
(V – R)	vs.	ν_c	97	$-0.36^{+0.12}_{-0.11}$	0.000303	3.61	24	0.002976	0.000648	0.38	0.43
(V – R)	vs.	i	97	$-0.36^{+0.12}_{-0.11}$	0.000323	3.60	25	0.003100	0.000675	0.38	0.43
(V – R)	vs.	ε	97	$-0.36^{+0.12}_{-0.11}$	0.000356	3.57	26	0.003224	0.000702	0.38	0.43
(B – I)	vs.	ν_c	53	$-0.47^{+0.14}_{-0.12}$	0.000357	3.57	27	0.003348	0.000729	0.51	0.57
(B – I)	vs.	ε	53	$-0.47^{+0.15}_{-0.12}$	0.000359	3.57	28	0.003472	0.000756	0.51	0.57
Grt	vs.	i	94	$-0.36^{+0.11}_{-0.10}$	0.000390	3.55	29	0.003596	0.000783	0.39	0.43
(B – I)	vs.	ψ	53	$-0.47^{+0.15}_{-0.13}$	0.000417	3.53	30	0.003720	0.000810	0.51	0.57
H_R	vs.	T_N	98	$0.34^{+0.09}_{-0.10}$	0.000507	3.48	31	0.003844	0.000837	0.38	0.43
(V – I)	vs.	(B – I)	53	$0.45^{+0.16}_{-0.20}$	0.000834	3.34	32	0.003968	0.000864	0.51	0.57
— 3 σ FDR correction cutoff —											
(V – R)	vs.	(V – I)	81	$0.36^{+0.13}_{-0.12}$	0.000940	3.31	33	0.004092	0.000891	0.42	0.47
(B – R)	vs.	q	79	$0.36^{+0.13}_{-0.13}$	0.001117	3.26	34	0.004216	0.000918	0.42	0.47
(B – V)	vs.	T_N	78	$0.36^{+0.13}_{-0.14}$	0.001192	3.24	35	0.004340	0.000945	0.42	0.47
H_R	vs.	i	98	$-0.32^{+0.10}_{-0.10}$	0.001426	3.19	36	0.004464	0.000972	0.38	0.43
H_R	vs.	ψ	98	$-0.32^{+0.10}_{-0.09}$	0.001585	3.16	37	0.004588	0.000999	0.38	0.43
H_R	vs.	ν_c	98	$-0.31^{+0.11}_{-0.10}$	0.001940	3.10	38	0.004712	0.001026	0.38	0.43
H_R	vs.	ε	98	$-0.30^{+0.10}_{-0.10}$	0.002265	3.05	39	0.004836	0.001053	0.38	0.43
(B – I)	vs.	i	53	$-0.41^{+0.16}_{-0.14}$	0.002328	3.04	40	0.004960	0.001080	0.51	0.57
(V – I)	vs.	(B – R)	65	$0.36^{+0.14}_{-0.16}$	0.003186	2.95	41	0.005084	0.001107	0.46	0.52
(R – I)	vs.	(V – I)	76	$0.33^{+0.15}_{-0.17}$	0.004019	2.88	42	0.005208	0.001134	0.43	0.48
— 2.5 σ FDR correction cutoff —											
Binary or multiple KBOs											
(B – I)	vs.	(B – R)	28	$0.82^{+0.09}_{-0.16}$	$< 10^{-6}$	5.42	1	0.000148	0.000032	0.67	0.73
(B – V)	vs.	(B – R)	34	$0.78^{+0.10}_{-0.16}$	$< 10^{-6}$	> 8.0	2	0.000295	0.000064	0.62	0.68
Grt	vs.	(B – R)	34	$0.74^{+0.10}_{-0.15}$	$< 10^{-6}$	5.06	3	0.000443	0.000096	0.62	0.68
Grt	vs.	(V – I)	37	$0.76^{+0.10}_{-0.16}$	$< 10^{-6}$	> 8.0	4	0.000590	0.000129	0.60	0.66
Grt	vs.	(V – R)	38	$0.76^{+0.10}_{-0.15}$	$< 10^{-6}$	> 8.0	5	0.000738	0.000161	0.59	0.65
(V – R)	vs.	(B – R)	34	$0.72^{+0.12}_{-0.18}$	0.000002	4.78	6	0.000886	0.000193	0.62	0.68
(V – I)	vs.	(B – R)	34	$0.70^{+0.11}_{-0.16}$	0.000004	4.60	7	0.001033	0.000225	0.62	0.68
(B – V)	vs.	(B – I)	28	$0.75^{+0.11}_{-0.18}$	0.000005	4.58	8	0.001181	0.000257	0.67	0.73
Grt	vs.	(B – I)	28	$0.75^{+0.11}_{-0.17}$	0.000005	4.57	9	0.001329	0.000289	0.67	0.73

continued on next page

Table 2. continued.

X^b	vs.	Y^c	n^d	$\rho_{-\sigma}^{+\sigma e}$	P-value ^f	$CL(\sigma)^g$	i^h	FDR correction ^a			
								$\frac{i}{m} 2.5 \sigma^i$	$\frac{i}{m} 3 \sigma^j$	$\rho_{\beta 2.5\sigma}^k$	$\rho_{\beta 3\sigma}^l$
(V – R)	vs.	(V – I)	37	0.67 ^{+0.12} _{-0.16}	0.000005	4.56	10	0.001476	0.000321	0.60	0.66
(V – I)	vs.	(B – I)	28	0.69 ^{+0.14} _{-0.24}	0.000045	4.08	11	0.001624	0.000354	0.67	0.73
(V – R)	vs.	(B – I)	28	0.69 ^{+0.12} _{-0.18}	0.000050	4.05	12	0.001771	0.000386	0.67	0.73
(B – R)	vs.	T_N	35	0.62 ^{+0.09} _{-0.12}	0.000063	4.00	13	0.001919	0.000418	0.61	0.68
(B – R)	vs.	v_c	35	-0.62 ^{+0.13} _{-0.11}	0.000067	3.98	14	0.002067	0.000450	0.61	0.68
Grt	vs.	T_N	38	0.56 ^{+0.12} _{-0.15}	0.000245	3.67	15	0.002214	0.000482	0.59	0.65
(B – R)	vs.	i	35	-0.58 ^{+0.15} _{-0.12}	0.000260	3.65	16	0.002362	0.000514	0.61	0.68
(B – I)	vs.	v_c	28	-0.63 ^{+0.16} _{-0.12}	0.000306	3.61	17	0.002510	0.000546	0.67	0.73
(B – V)	vs.	(V – I)	33	0.59 ^{+0.14} _{-0.12}	0.000331	3.59	18	0.002657	0.000579	0.63	0.69
Grt	vs.	(B – V)	34	0.57 ^{+0.15} _{-0.13}	0.000410	3.53	19	0.002805	0.000611	0.62	0.68
(V – I)	vs.	T_N	47	0.49 ^{+0.13} _{-0.16}	0.000446	3.51	20	0.002952	0.000643	0.54	0.60
— 3 σ FDR correction cutoff —											
(V – R)	vs.	T_N	38	0.53 ^{+0.13} _{-0.16}	0.000706	3.39	21	0.003100	0.000675	0.59	0.65
(B – V)	vs.	v_c	34	-0.55 ^{+0.16} _{-0.13}	0.000750	3.37	22	0.003248	0.000707	0.62	0.68
(B – V)	vs.	(V – R)	34	0.55 ^{+0.15} _{-0.20}	0.000815	3.35	23	0.003395	0.000739	0.62	0.68
(B – I)	vs.	T_N	28	0.59 ^{+0.13} _{-0.17}	0.000858	3.33	24	0.003543	0.000771	0.67	0.73
Grt	vs.	v_c	38	-0.51 ^{+0.15} _{-0.13}	0.001201	3.24	25	0.003690	0.000804	0.59	0.65
(B – V)	vs.	T_N	34	0.52 ^{+0.12} _{-0.14}	0.001522	3.17	26	0.003838	0.000836	0.62	0.68
Grt	vs.	(R – I)	35	0.50 ^{+0.17} _{-0.22}	0.002226	3.06	27	0.003986	0.000868	0.61	0.68
H_R	vs.	i	38	-0.48 ^{+0.15} _{-0.13}	0.002353	3.04	28	0.004133	0.000900	0.59	0.65
(V – R)	vs.	v_c	38	-0.48 ^{+0.17} _{-0.14}	0.002437	3.03	29	0.004281	0.000932	0.59	0.65
(V – I)	vs.	v_c	47	-0.43 ^{+0.17} _{-0.15}	0.002467	3.03	30	0.004429	0.000964	0.54	0.60
(R – I)	vs.	(V – I)	35	0.49 ^{+0.20} _{-0.27}	0.002609	3.01	31	0.004576	0.000996	0.61	0.68
Grt	vs.	i	38	-0.47 ^{+0.16} _{-0.13}	0.003044	2.96	32	0.004724	0.001029	0.59	0.65
(V – R)	vs.	i	38	-0.46 ^{+0.17} _{-0.14}	0.003376	2.93	33	0.004871	0.001061	0.59	0.65
(B – V)	vs.	i	34	-0.48 ^{+0.17} _{-0.14}	0.003660	2.91	34	0.005019	0.001093	0.62	0.68
(B – I)	vs.	i	28	-0.53 ^{+0.19} _{-0.15}	0.003746	2.90	35	0.005167	0.001125	0.67	0.73
— 2.5 σ FDR correction cutoff —											
KBO*											
(B – I)	vs.	(B – R)	207	0.83 ^{+0.04} _{-0.05}	< 10 ⁻⁶	> 8.0	1	0.000148	0.000032	0.26	0.30
(B – I)	vs.	T_N	209	0.36 ^{+0.07} _{-0.07}	< 10 ⁻⁶	5.42	2	0.000295	0.000064	0.26	0.29
(B – I)	vs.	v_c	209	-0.39 ^{+0.07} _{-0.06}	< 10 ⁻⁶	> 8.0	3	0.000443	0.000096	0.26	0.29
(B – R)	vs.	T_N	268	0.41 ^{+0.06} _{-0.06}	< 10 ⁻⁶	> 8.0	4	0.000590	0.000129	0.23	0.26
(B – R)	vs.	v_c	268	-0.42 ^{+0.06} _{-0.06}	< 10 ⁻⁶	> 8.0	5	0.000738	0.000161	0.23	0.26
(B – R)	vs.	i	268	-0.39 ^{+0.06} _{-0.05}	< 10 ⁻⁶	> 8.0	6	0.000886	0.000193	0.23	0.26
(B – V)	vs.	(B – I)	209	0.75 ^{+0.05} _{-0.05}	< 10 ⁻⁶	> 8.0	7	0.001033	0.000225	0.26	0.29
(B – V)	vs.	(B – R)	264	0.76 ^{+0.04} _{-0.05}	< 10 ⁻⁶	> 8.0	8	0.001181	0.000257	0.23	0.26
(B – V)	vs.	(R – I)	223	0.41 ^{+0.07} _{-0.08}	< 10 ⁻⁶	> 8.0	9	0.001329	0.000289	0.25	0.29
(B – V)	vs.	T_N	266	0.35 ^{+0.06} _{-0.06}	< 10 ⁻⁶	> 8.0	10	0.001476	0.000321	0.23	0.26
(B – V)	vs.	(V – I)	233	0.58 ^{+0.06} _{-0.06}	< 10 ⁻⁶	> 8.0	11	0.001624	0.000354	0.25	0.28
(B – V)	vs.	(V – R)	266	0.51 ^{+0.07} _{-0.07}	< 10 ⁻⁶	> 8.0	12	0.001771	0.000386	0.23	0.26
(B – V)	vs.	v_c	266	-0.37 ^{+0.06} _{-0.06}	< 10 ⁻⁶	> 8.0	13	0.001919	0.000418	0.23	0.26
(B – V)	vs.	i	266	-0.34 ^{+0.06} _{-0.06}	< 10 ⁻⁶	> 8.0	14	0.002067	0.000450	0.23	0.26
Grt	vs.	(B – I)	209	0.77 ^{+0.05} _{-0.06}	< 10 ⁻⁶	> 8.0	15	0.002214	0.000482	0.26	0.29
Grt	vs.	(B – R)	264	0.74 ^{+0.04} _{-0.05}	< 10 ⁻⁶	> 8.0	16	0.002362	0.000514	0.23	0.26
Grt	vs.	(B – V)	266	0.59 ^{+0.06} _{-0.06}	< 10 ⁻⁶	> 8.0	17	0.002510	0.000546	0.23	0.26
Grt	vs.	(R – I)	251	0.48 ^{+0.07} _{-0.07}	< 10 ⁻⁶	> 8.0	18	0.002657	0.000579	0.24	0.27
Grt	vs.	T_N	294	0.36 ^{+0.06} _{-0.06}	< 10 ⁻⁶	> 8.0	19	0.002805	0.000611	0.22	0.25
Grt	vs.	(V – I)	261	0.72 ^{+0.04} _{-0.05}	< 10 ⁻⁶	> 8.0	20	0.002952	0.000643	0.23	0.26
Grt	vs.	(V – R)	294	0.71 ^{+0.05} _{-0.06}	< 10 ⁻⁶	> 8.0	21	0.003100	0.000675	0.22	0.25
Grt	vs.	v_c	294	-0.36 ^{+0.06} _{-0.06}	< 10 ⁻⁶	> 8.0	22	0.003248	0.000707	0.22	0.25
Grt	vs.	i	294	-0.35 ^{+0.06} _{-0.06}	< 10 ⁻⁶	> 8.0	23	0.003395	0.000739	0.22	0.25
H_R	vs.	q	291	-0.37 ^{+0.05} _{-0.05}	< 10 ⁻⁶	> 8.0	24	0.003543	0.000771	0.22	0.25

continued on next page

Table 2. continued.

X^b	vs.	Y^c	n^d	$\rho_{-\sigma}^{+ \sigma e}$	P-value ^f	$CL(\sigma)^g$	i^h	FDR correction ^a			
								$\frac{i}{m} 2.5 \sigma^i$	$\frac{i}{m} 3 \sigma^j$	$\rho_{\beta 2.5 \sigma}^k$	$\rho_{\beta 3 \sigma}^l$
(R – I)	vs.	(B – I)	206	0.56 ^{+0.07} _{-0.07}	< 10 ⁻⁶	> 8.0	25	0.003690	0.000804	0.26	0.30
(R – I)	vs.	(B – R)	222	0.44 ^{+0.07} _{-0.08}	< 10 ⁻⁶	> 8.0	26	0.003838	0.000836	0.25	0.29
(R – I)	vs.	(V – I)	253	0.53 ^{+0.07} _{-0.08}	< 10 ⁻⁶	> 8.0	27	0.003986	0.000868	0.24	0.27
(V – I)	vs.	(B – I)	209	0.73 ^{+0.05} _{-0.06}	< 10 ⁻⁶	> 8.0	28	0.004133	0.000900	0.26	0.29
(V – I)	vs.	(B – R)	232	0.68 ^{+0.05} _{-0.06}	< 10 ⁻⁶	> 8.0	29	0.004281	0.000932	0.25	0.28
(V – I)	vs.	T_N	281	0.33 ^{+0.06} _{-0.06}	< 10 ⁻⁶	> 8.0	30	0.004429	0.000964	0.23	0.25
(V – I)	vs.	v_c	281	-0.34 ^{+0.06} _{-0.06}	< 10 ⁻⁶	> 8.0	31	0.004576	0.000996	0.23	0.25
(V – I)	vs.	i	281	-0.31 ^{+0.06} _{-0.06}	< 10 ⁻⁶	5.42	32	0.004724	0.001029	0.23	0.25
(V – R)	vs.	(B – I)	209	0.65 ^{+0.06} _{-0.08}	< 10 ⁻⁶	> 8.0	33	0.004871	0.001061	0.26	0.29
(V – R)	vs.	(B – R)	264	0.67 ^{+0.06} _{-0.07}	< 10 ⁻⁶	> 8.0	34	0.005019	0.001093	0.23	0.26
(V – R)	vs.	T_N	297	0.32 ^{+0.06} _{-0.07}	< 10 ⁻⁶	> 8.0	35	0.005167	0.001125	0.22	0.25
(V – R)	vs.	(V – I)	262	0.59 ^{+0.05} _{-0.06}	< 10 ⁻⁶	> 8.0	36	0.005314	0.001157	0.23	0.26
(V – R)	vs.	v_c	297	-0.32 ^{+0.07} _{-0.06}	< 10 ⁻⁶	> 8.0	37	0.005462	0.001189	0.22	0.25
(V – R)	vs.	i	297	-0.32 ^{+0.07} _{-0.06}	< 10 ⁻⁶	> 8.0	38	0.005610	0.001221	0.22	0.25
(B – I)	vs.	i	209	-0.33 ^{+0.07} _{-0.07}	0.000001	4.89	39	0.005757	0.001254	0.26	0.29
(V – R)	vs.	(R – I)	252	0.30 ^{+0.08} _{-0.08}	0.000001	4.83	40	0.005905	0.001286	0.24	0.27
(B – R)	vs.	e	268	-0.25 ^{+0.06} _{-0.06}	0.000035	4.14	41	0.006052	0.001318	0.23	0.26
(B – R)	vs.	q	268	0.25 ^{+0.06} _{-0.06}	0.000038	4.12	42	0.006200	0.001350	0.23	0.26
Grt	vs.	q	294	0.22 ^{+0.07} _{-0.07}	0.000170	3.76	43	0.006348	0.001382	0.22	0.25
Grt	vs.	e	294	-0.21 ^{+0.06} _{-0.06}	0.000259	3.65	44	0.006495	0.001414	0.22	0.25
(B – V)	vs.	q	266	0.22 ^{+0.06} _{-0.07}	0.000313	3.60	45	0.006643	0.001446	0.23	0.26
(B – I)	vs.	q	209	0.24 ^{+0.07} _{-0.07}	0.000447	3.51	46	0.006790	0.001479	0.26	0.29
(B – V)	vs.	e	266	-0.21 ^{+0.06} _{-0.06}	0.000466	3.50	47	0.006938	0.001511	0.23	0.26
(B – I)	vs.	e	209	-0.24 ^{+0.07} _{-0.06}	0.000470	3.50	48	0.007086	0.001543	0.26	0.29
(V – I)	vs.	e	281	-0.21 ^{+0.06} _{-0.06}	0.000473	3.50	49	0.007233	0.001575	0.23	0.25
(V – I)	vs.	q	281	0.20 ^{+0.07} _{-0.07}	0.000588	3.44	50	0.007381	0.001607	0.23	0.25
— 3 σ FDR correction cutoff —											
(R – I)	vs.	v_c	253	-0.20 ^{+0.08} _{-0.07}	0.001711	3.14	51	0.007529	0.001639	0.24	0.27
(R – I)	vs.	T_N	253	0.19 ^{+0.07} _{-0.08}	0.002341	3.04	52	0.007676	0.001671	0.24	0.27
H_R	vs.	a	291	-0.18 ^{+0.06} _{-0.06}	0.002384	3.04	53	0.007824	0.001704	0.22	0.25
(V – R)	vs.	q	297	0.17 ^{+0.06} _{-0.07}	0.004010	2.88	54	0.007971	0.001736	0.22	0.25
(V – R)	vs.	e	297	-0.16 ^{+0.07} _{-0.06}	0.004778	2.82	55	0.008119	0.001768	0.22	0.25
H_R	vs.	e	291	0.16 ^{+0.06} _{-0.06}	0.005972	2.75	56	0.008267	0.001800	0.22	0.25
(R – I)	vs.	q	253	0.17 ^{+0.07} _{-0.08}	0.007545	2.67	57	0.008414	0.001832	0.24	0.27
— 2.5 σ FDR correction cutoff —											
All*											
(B – I)	vs.	(B – R)	226	0.84 ^{+0.04} _{-0.05}	< 10 ⁻⁶	> 8.0	1	0.000148	0.000032	0.25	0.28
(B – I)	vs.	v_c	228	-0.36 ^{+0.07} _{-0.06}	< 10 ⁻⁶	> 8.0	2	0.000295	0.000064	0.25	0.28
(B – I)	vs.	i	228	-0.33 ^{+0.06} _{-0.06}	< 10 ⁻⁶	5.06	3	0.000443	0.000096	0.25	0.28
(B – R)	vs.	T_N	297	0.29 ^{+0.06} _{-0.06}	< 10 ⁻⁶	5.22	4	0.000590	0.000129	0.22	0.25
(B – R)	vs.	v_c	297	-0.41 ^{+0.06} _{-0.05}	< 10 ⁻⁶	> 8.0	5	0.000738	0.000161	0.22	0.25
(B – R)	vs.	i	297	-0.39 ^{+0.06} _{-0.05}	< 10 ⁻⁶	> 8.0	6	0.000886	0.000193	0.22	0.25
(B – V)	vs.	(B – I)	228	0.77 ^{+0.04} _{-0.05}	< 10 ⁻⁶	> 8.0	7	0.001033	0.000225	0.25	0.28
(B – V)	vs.	(B – R)	290	0.78 ^{+0.04} _{-0.04}	< 10 ⁻⁶	> 8.0	8	0.001181	0.000257	0.22	0.25
(B – V)	vs.	(R – I)	246	0.44 ^{+0.07} _{-0.07}	< 10 ⁻⁶	> 8.0	9	0.001329	0.000289	0.24	0.27
(B – V)	vs.	(V – I)	256	0.61 ^{+0.05} _{-0.06}	< 10 ⁻⁶	> 8.0	10	0.001476	0.000321	0.24	0.27
(B – V)	vs.	(V – R)	292	0.54 ^{+0.06} _{-0.07}	< 10 ⁻⁶	> 8.0	11	0.001624	0.000354	0.22	0.25
(B – V)	vs.	v_c	292	-0.37 ^{+0.06} _{-0.06}	< 10 ⁻⁶	> 8.0	12	0.001771	0.000386	0.22	0.25
(B – V)	vs.	i	292	-0.35 ^{+0.06} _{-0.06}	< 10 ⁻⁶	> 8.0	13	0.001919	0.000418	0.22	0.25
Grt	vs.	(B – I)	228	0.78 ^{+0.04} _{-0.05}	< 10 ⁻⁶	> 8.0	14	0.002067	0.000450	0.25	0.28
Grt	vs.	(B – R)	292	0.76 ^{+0.04} _{-0.05}	< 10 ⁻⁶	> 8.0	15	0.002214	0.000482	0.22	0.25
Grt	vs.	(B – V)	292	0.61 ^{+0.05} _{-0.06}	< 10 ⁻⁶	> 8.0	16	0.002362	0.000514	0.22	0.25
Grt	vs.	(R – I)	279	0.52 ^{+0.06} _{-0.07}	< 10 ⁻⁶	> 8.0	17	0.002510	0.000546	0.23	0.26

continued on next page

Table 2. continued.

X^b	vs.	Y^c	n^d	$\rho_{-\sigma}^{+\sigma e}$	P-value ^f	$CL(\sigma)^g$	FDR correction ^a				
							i^h	$\frac{i}{m} 2.5 \sigma^i$	$\frac{i}{m} 3 \sigma^j$	$\rho_{\beta 2.5\sigma}^k$	$\rho_{\beta 3\sigma}^l$
<i>Grt</i>	vs.	T_N	325	$0.28^{+0.06}_{-0.06}$	$< 10^{-6}$	5.12	18	0.002657	0.000579	0.21	0.24
<i>Grt</i>	vs.	$(V - I)$	289	$0.74^{+0.04}_{-0.04}$	$< 10^{-6}$	> 8.0	19	0.002805	0.000611	0.22	0.25
<i>Grt</i>	vs.	$(V - R)$	325	$0.72^{+0.04}_{-0.05}$	$< 10^{-6}$	> 8.0	20	0.002952	0.000643	0.21	0.24
<i>Grt</i>	vs.	v_c	325	$-0.33^{+0.06}_{-0.06}$	$< 10^{-6}$	> 8.0	21	0.003100	0.000675	0.21	0.24
<i>Grt</i>	vs.	i	325	$-0.34^{+0.06}_{-0.05}$	$< 10^{-6}$	> 8.0	22	0.003248	0.000707	0.21	0.24
H_R	vs.	a	323	$-0.36^{+0.06}_{-0.06}$	$< 10^{-6}$	> 8.0	23	0.003395	0.000739	0.21	0.24
H_R	vs.	q	323	$-0.51^{+0.05}_{-0.04}$	$< 10^{-6}$	> 8.0	24	0.003543	0.000771	0.21	0.24
$(R - I)$	vs.	$(B - I)$	225	$0.58^{+0.06}_{-0.07}$	$< 10^{-6}$	> 8.0	25	0.003690	0.000804	0.25	0.28
$(R - I)$	vs.	$(B - R)$	247	$0.47^{+0.07}_{-0.07}$	$< 10^{-6}$	> 8.0	26	0.003838	0.000836	0.24	0.27
$(R - I)$	vs.	$(V - I)$	281	$0.56^{+0.06}_{-0.07}$	$< 10^{-6}$	> 8.0	27	0.003986	0.000868	0.23	0.25
$(V - I)$	vs.	$(B - I)$	228	$0.75^{+0.05}_{-0.06}$	$< 10^{-6}$	> 8.0	28	0.004133	0.000900	0.25	0.28
$(V - I)$	vs.	$(B - R)$	257	$0.70^{+0.04}_{-0.05}$	$< 10^{-6}$	> 8.0	29	0.004281	0.000932	0.24	0.27
$(V - I)$	vs.	v_c	309	$-0.31^{+0.06}_{-0.06}$	$< 10^{-6}$	> 8.0	30	0.004429	0.000964	0.21	0.24
$(V - I)$	vs.	i	309	$-0.30^{+0.06}_{-0.06}$	$< 10^{-6}$	5.42	31	0.004576	0.000996	0.21	0.24
$(V - R)$	vs.	$(B - I)$	228	$0.66^{+0.06}_{-0.07}$	$< 10^{-6}$	> 8.0	32	0.004724	0.001029	0.25	0.28
$(V - R)$	vs.	$(B - R)$	292	$0.68^{+0.05}_{-0.06}$	$< 10^{-6}$	> 8.0	33	0.004871	0.001061	0.22	0.25
$(V - R)$	vs.	$(R - I)$	280	$0.34^{+0.07}_{-0.08}$	$< 10^{-6}$	> 8.0	34	0.005019	0.001093	0.23	0.25
$(V - R)$	vs.	$(V - I)$	290	$0.61^{+0.05}_{-0.06}$	$< 10^{-6}$	> 8.0	35	0.005167	0.001125	0.22	0.25
$(V - R)$	vs.	v_c	328	$-0.29^{+0.06}_{-0.06}$	$< 10^{-6}$	5.42	36	0.005314	0.001157	0.21	0.24
$(V - R)$	vs.	i	328	$-0.31^{+0.06}_{-0.06}$	$< 10^{-6}$	> 8.0	37	0.005462	0.001189	0.21	0.24
$(B - V)$	vs.	T_N	292	$0.28^{+0.06}_{-0.07}$	0.000001	4.85	38	0.005610	0.001221	0.22	0.25
$(V - I)$	vs.	T_N	309	$0.26^{+0.07}_{-0.07}$	0.000005	4.56	39	0.005757	0.001254	0.21	0.24
$(B - R)$	vs.	q	297	$0.26^{+0.06}_{-0.06}$	0.000006	4.52	40	0.005905	0.001286	0.22	0.25
H_R	vs.	e	323	$0.25^{+0.05}_{-0.05}$	0.000006	4.54	41	0.006052	0.001318	0.21	0.24
$(V - R)$	vs.	T_N	328	$0.24^{+0.06}_{-0.07}$	0.000008	4.47	42	0.006200	0.001350	0.21	0.24
H_R	vs.	Q	323	$-0.24^{+0.06}_{-0.06}$	0.000016	4.31	43	0.006348	0.001382	0.21	0.24
$(B - R)$	vs.	e	297	$-0.24^{+0.06}_{-0.05}$	0.000038	4.12	44	0.006495	0.001414	0.22	0.25
$(B - V)$	vs.	q	292	$0.24^{+0.06}_{-0.06}$	0.000040	4.11	45	0.006643	0.001446	0.22	0.25
$(B - I)$	vs.	T_N	228	$0.26^{+0.07}_{-0.07}$	0.000056	4.03	46	0.006790	0.001479	0.25	0.28
H_R	vs.	v_c	323	$0.21^{+0.06}_{-0.06}$	0.000117	3.85	47	0.006938	0.001511	0.21	0.24
$(B - I)$	vs.	q	228	$0.25^{+0.07}_{-0.07}$	0.000163	3.77	48	0.007086	0.001543	0.25	0.28
$(B - V)$	vs.	e	292	$-0.21^{+0.06}_{-0.06}$	0.000294	3.62	49	0.007233	0.001575	0.22	0.25
$(V - I)$	vs.	q	309	$0.20^{+0.06}_{-0.06}$	0.000329	3.59	50	0.007381	0.001607	0.21	0.24
<i>Grt</i>	vs.	e	325	$-0.20^{+0.06}_{-0.06}$	0.000350	3.58	51	0.007529	0.001639	0.21	0.24
<i>Grt</i>	vs.	q	325	$0.20^{+0.06}_{-0.06}$	0.000399	3.54	52	0.007676	0.001671	0.21	0.24
$(V - I)$	vs.	e	309	$-0.19^{+0.06}_{-0.06}$	0.000667	3.40	53	0.007824	0.001704	0.21	0.24
$(B - I)$	vs.	e	228	$-0.21^{+0.06}_{-0.06}$	0.001104	3.26	54	0.007971	0.001736	0.25	0.28
$(R - I)$	vs.	v_c	281	$-0.19^{+0.07}_{-0.07}$	0.001697	3.14	55	0.008119	0.001768	0.23	0.25
— 3 σ FDR correction cutoff —											
H_R	vs.	$(B - R)$	281	$-0.17^{+0.07}_{-0.07}$	0.004567	2.84	56	0.008267	0.001800	0.23	0.25
$(R - I)$	vs.	i	281	$-0.17^{+0.07}_{-0.07}$	0.005277	2.79	57	0.008414	0.001832	0.23	0.25
$(V - R)$	vs.	e	328	$-0.15^{+0.06}_{-0.06}$	0.005446	2.78	58	0.008562	0.001864	0.21	0.24
$(V - R)$	vs.	q	328	$0.15^{+0.06}_{-0.06}$	0.006765	2.71	59	0.008710	0.001896	0.21	0.24
$(R - I)$	vs.	q	281	$0.16^{+0.07}_{-0.07}$	0.007999	2.65	60	0.008857	0.001929	0.23	0.25
— 2.5 σ FDR correction cutoff —											
KBOS*exceptforCKBOs											
$(B - I)$	vs.	$(B - R)$	154	$0.86^{+0.04}_{-0.06}$	$< 10^{-6}$	> 8.0	1	0.000148	0.000032	0.30	0.34
$(B - V)$	vs.	$(B - I)$	156	$0.79^{+0.05}_{-0.06}$	$< 10^{-6}$	> 8.0	2	0.000295	0.000064	0.30	0.34
$(B - V)$	vs.	$(B - R)$	183	$0.81^{+0.04}_{-0.05}$	$< 10^{-6}$	> 8.0	3	0.000443	0.000096	0.28	0.31
$(B - V)$	vs.	$(R - I)$	163	$0.46^{+0.08}_{-0.09}$	$< 10^{-6}$	> 8.0	4	0.000590	0.000129	0.29	0.33
$(B - V)$	vs.	$(V - I)$	168	$0.64^{+0.06}_{-0.07}$	$< 10^{-6}$	> 8.0	5	0.000738	0.000161	0.29	0.33
$(B - V)$	vs.	$(V - R)$	185	$0.61^{+0.07}_{-0.08}$	$< 10^{-6}$	> 8.0	6	0.000886	0.000193	0.28	0.31
<i>Grt</i>	vs.	$(B - I)$	156	$0.81^{+0.06}_{-0.07}$	$< 10^{-6}$	> 8.0	7	0.001033	0.000225	0.30	0.34

continued on next page

Table 2. continued.

X^b	vs.	Y^c	n^d	$\rho_{-\sigma}^{+ \sigma e}$	P-value ^f	$CL(\sigma)^g$	FDR correction ^a				
							i^h	$\frac{i}{m} 2.5 \sigma^i$	$\frac{i}{m} 3 \sigma^j$	$\rho_{\beta 2.5 \sigma}^k$	$\rho_{\beta 3 \sigma}^l$
<i>Grt</i>	vs.	(<i>B</i> – <i>R</i>)	183	$0.79^{+0.05}_{-0.07}$	$< 10^{-6}$	> 8.0	8	0.001181	0.000257	0.28	0.31
<i>Grt</i>	vs.	(<i>B</i> – <i>V</i>)	185	$0.68^{+0.06}_{-0.07}$	$< 10^{-6}$	> 8.0	9	0.001329	0.000289	0.28	0.31
<i>Grt</i>	vs.	(<i>R</i> – <i>I</i>)	175	$0.53^{+0.08}_{-0.09}$	$< 10^{-6}$	> 8.0	10	0.001476	0.000321	0.28	0.32
<i>Grt</i>	vs.	(<i>V</i> – <i>I</i>)	180	$0.76^{+0.05}_{-0.06}$	$< 10^{-6}$	> 8.0	11	0.001624	0.000354	0.28	0.32
<i>Grt</i>	vs.	(<i>V</i> – <i>R</i>)	197	$0.75^{+0.05}_{-0.06}$	$< 10^{-6}$	> 8.0	12	0.001771	0.000386	0.27	0.30
<i>H_R</i>	vs.	<i>q</i>	191	$-0.39^{+0.07}_{-0.06}$	$< 10^{-6}$	> 8.0	13	0.001919	0.000418	0.27	0.31
(<i>R</i> – <i>I</i>)	vs.	(<i>B</i> – <i>I</i>)	154	$0.62^{+0.07}_{-0.09}$	$< 10^{-6}$	> 8.0	14	0.002067	0.000450	0.30	0.34
(<i>R</i> – <i>I</i>)	vs.	(<i>B</i> – <i>R</i>)	162	$0.49^{+0.08}_{-0.09}$	$< 10^{-6}$	> 8.0	15	0.002214	0.000482	0.30	0.33
(<i>R</i> – <i>I</i>)	vs.	(<i>V</i> – <i>I</i>)	176	$0.59^{+0.07}_{-0.08}$	$< 10^{-6}$	> 8.0	16	0.002362	0.000514	0.28	0.32
(<i>V</i> – <i>I</i>)	vs.	(<i>B</i> – <i>I</i>)	156	$0.78^{+0.05}_{-0.07}$	$< 10^{-6}$	> 8.0	17	0.002510	0.000546	0.30	0.34
(<i>V</i> – <i>I</i>)	vs.	(<i>B</i> – <i>R</i>)	166	$0.72^{+0.05}_{-0.06}$	$< 10^{-6}$	> 8.0	18	0.002657	0.000579	0.29	0.33
(<i>V</i> – <i>R</i>)	vs.	(<i>B</i> – <i>I</i>)	156	$0.69^{+0.08}_{-0.09}$	$< 10^{-6}$	> 8.0	19	0.002805	0.000611	0.30	0.34
(<i>V</i> – <i>R</i>)	vs.	(<i>B</i> – <i>R</i>)	183	$0.72^{+0.07}_{-0.09}$	$< 10^{-6}$	> 8.0	20	0.002952	0.000643	0.28	0.31
(<i>V</i> – <i>R</i>)	vs.	(<i>V</i> – <i>I</i>)	180	$0.63^{+0.07}_{-0.08}$	$< 10^{-6}$	> 8.0	21	0.003100	0.000675	0.28	0.32
(<i>V</i> – <i>R</i>)	vs.	(<i>R</i> – <i>I</i>)	175	$0.36^{+0.09}_{-0.10}$	0.000001	4.85	22	0.003248	0.000707	0.28	0.32
(<i>B</i> – <i>R</i>)	vs.	<i>v_c</i>	186	$-0.30^{+0.07}_{-0.07}$	0.000030	4.17	23	0.003395	0.000739	0.28	0.31
(<i>B</i> – <i>R</i>)	vs.	<i>i</i>	186	$-0.30^{+0.07}_{-0.07}$	0.000042	4.09	24	0.003543	0.000771	0.28	0.31
(<i>B</i> – <i>R</i>)	vs.	<i>T_N</i>	186	$0.29^{+0.07}_{-0.07}$	0.000076	3.96	25	0.003690	0.000804	0.28	0.31
<i>Grt</i>	vs.	<i>i</i>	197	$-0.28^{+0.07}_{-0.07}$	0.000076	3.96	26	0.003838	0.000836	0.27	0.30
(<i>B</i> – <i>V</i>)	vs.	<i>i</i>	185	$-0.27^{+0.08}_{-0.08}$	0.000194	3.73	27	0.003986	0.000868	0.28	0.31
(<i>B</i> – <i>V</i>)	vs.	<i>v_c</i>	185	$-0.27^{+0.08}_{-0.08}$	0.000200	3.72	28	0.004133	0.000900	0.28	0.31
(<i>V</i> – <i>R</i>)	vs.	<i>i</i>	197	$-0.26^{+0.08}_{-0.08}$	0.000262	3.65	29	0.004281	0.000932	0.27	0.30
(<i>B</i> – <i>V</i>)	vs.	<i>T_N</i>	185	$0.26^{+0.08}_{-0.08}$	0.000271	3.64	30	0.004429	0.000964	0.28	0.31
<i>Grt</i>	vs.	<i>v_c</i>	197	$-0.25^{+0.08}_{-0.07}$	0.000468	3.50	31	0.004576	0.000996	0.27	0.30
<i>Grt</i>	vs.	<i>T_N</i>	197	$0.24^{+0.07}_{-0.07}$	0.000792	3.36	32	0.004724	0.001029	0.27	0.30
— 3 σ FDR correction cutoff —											
(<i>B</i> – <i>I</i>)	vs.	<i>T_N</i>	156	$0.27^{+0.08}_{-0.08}$	0.000796	3.35	33	0.004871	0.001061	0.30	0.34
(<i>B</i> – <i>I</i>)	vs.	<i>v_c</i>	156	$-0.27^{+0.08}_{-0.08}$	0.000804	3.35	34	0.005019	0.001093	0.30	0.34
(<i>V</i> – <i>R</i>)	vs.	<i>v_c</i>	197	$-0.23^{+0.08}_{-0.08}$	0.001228	3.23	35	0.005167	0.001125	0.27	0.30
(<i>V</i> – <i>I</i>)	vs.	<i>i</i>	182	$-0.24^{+0.08}_{-0.08}$	0.001251	3.23	36	0.005314	0.001157	0.28	0.32
(<i>V</i> – <i>I</i>)	vs.	<i>v_c</i>	182	$-0.24^{+0.08}_{-0.08}$	0.001263	3.22	37	0.005462	0.001189	0.28	0.32
(<i>B</i> – <i>I</i>)	vs.	<i>i</i>	156	$-0.25^{+0.08}_{-0.08}$	0.001330	3.21	38	0.005610	0.001221	0.30	0.34
(<i>V</i> – <i>I</i>)	vs.	<i>T_N</i>	182	$0.22^{+0.08}_{-0.08}$	0.002389	3.04	39	0.005757	0.001254	0.28	0.32
(<i>V</i> – <i>R</i>)	vs.	<i>T_N</i>	197	$0.21^{+0.08}_{-0.08}$	0.002943	2.97	40	0.005905	0.001286	0.27	0.30
<i>H_R</i>	vs.	<i>a</i>	191	$-0.19^{+0.07}_{-0.07}$	0.007272	2.68	41	0.006052	0.001318	0.27	0.31
<i>H_R</i>	vs.	(<i>B</i> – <i>R</i>)	176	$-0.16^{+0.08}_{-0.08}$	0.028693	2.19	42	0.006200	0.001350	0.28	0.32
— 2.5 σ FDR correction cutoff —											

^(a) Computational steps of the correction for the false-discovery rate (see Sect. 3.4) ^(b) X variable. ^(c) Y variable. ^(d) Number of data points.

^(e) Spearman- ρ correlation value $\pm$ the 68.2% uncertainty. ^(f) P-value of the correlation.

^(g) Confidence level of the correlation in σ values of a normalized Gaussian distribution.

^(h) Rank of the correlation after ordering by increasing P-values

⁽ⁱ⁾ FDR-corrected P-value as a function of the number of tests for 2.5 σ confidence.

^(j) FDR-corrected P-value as a function of the number of tests for 3 σ confidence.

^(k) Maximum detectable ρ at a 2.5 σ confidence with a β risk of failing to detect it of 10%.

^(l) Maximum detectable ρ at a 3 σ confidence with a β risk of failing to detect it of 10%.

Table 5. Compilation of absolute magnitude and visible colors and corresponding orbital parameters

Object	Class ^a	H_R^b	GrI^c	$B - V$	$V - R$	$R - I$	$V - I$	$B - I$	$B - R$	q	Q	i	e	a	u_c^d	u_f^d	e_f^d	T_N^g	References	
1993RO	Plut	8.48 ± 0.11	19.02 ± 8.66	0.86 ± 0.11	0.58 ± 0.13	0.52 ± 0.19	1.02 ± 0.22	1.70 ± 0.29	1.39 ± 0.15	31.521	47.275	3.7	0.1999	39.398	1.00	0.0007406	0.210	3.002	5.513 1, 2, 3	
1994ES ₂	cClas	7.53 ± 0.12	74.75 ± 9.01	—	—	—	—	—	—	—	50.977	1.1	0.1123	45.832	0.50	0.0001795	0.114	3.109	6.011 1	
1994EV ₃	cClas	7.12 ± 0.07	29.48 ± 7.07	1.09 ± 0.16	0.58 ± 0.12	0.80 ± 0.10	1.47 ± 0.18	2.54 ± 0.21	1.73 ± 0.17	40.903	44.690	1.7	0.0442	42.797	0.27	0.0000481	0.053	3.085	5.849 1, 4, 5, 6	
1994TA	Cent	11.40 ± 0.13	32.61 ± 6.24	1.26 ± 0.14	0.67 ± 0.08	0.74 ± 0.21	1.36 ± 0.29	2.72 ± 0.41	1.93 ± 0.15	11.791	21.830	5.4	0.2986	16.811	2.24	0.0038409	0.313	3.210	3.725 3, 7	
1995DB ₂	cClas	8.11 ± 0.09	—	—	—	—	—	—	—	—	53.041	4.1	0.1383	46.596	0.68	0.0003663	0.156	3.105	6.024 8	
1995DC ₂	cClas	6.79 ± 0.16	42.22 ± 10.48	—	—	—	—	—	—	—	46.798	2.3	0.0600	44.149	0.33	0.0000888	0.073	3.098	5.928 9	
1995FB ₂₁	cClas	7.02 ± 0.10	—	—	—	—	—	—	—	—	50.896	0.7	0.1144	45.671	0.51	0.0001822	0.115	3.107	6.000 9	
1995HM ₅	Plut	7.86 ± 0.11	8.38 ± 6.31	0.60 ± 0.15	0.46 ± 0.10	0.43 ± 0.13	0.93 ± 0.10	—	—	—	49.252	4.8	0.2497	39.410	1.25	0.0011680	0.263	2.972	5.443 4, 8, 9, 10	
1995WY ₂	cClas	6.86 ± 0.11	22.71 ± 11.70	1.00 ± 0.21	0.65 ± 0.19	0.46 ± 0.21	1.11 ± 0.28	2.12 ± 0.34	1.66 ± 0.28	40.814	52.677	1.6	0.1269	46.745	0.57	0.0002329	0.130	3.116	6.055 1, 7	
1996RQ ₂₀	hClas	6.90 ± 0.09	18.92 ± 6.22	0.93 ± 0.14	0.54 ± 0.10	0.61 ± 0.11	1.20 ± 0.12	2.16 ± 0.17	1.52 ± 0.16	39.492	48.605	31.7	0.1034	44.049	2.52	0.0064047	0.535	2.732	5.845 7, 2, 8, 9	
1996RR ₂₀	Plut	6.62 ± 0.14	35.42 ± 6.21	1.14 ± 0.11	0.73 ± 0.10	0.63 ± 0.12	1.39 ± 0.17	2.50 ± 0.24	1.87 ± 0.13	32.720	46.413	5.3	0.1730	39.567	0.93	0.0006891	0.196	3.010	5.540 7, 3, 5, 8	
1996TC ₆₈	hClas	6.73 ± 0.07	—	—	—	—	—	—	—	—	38.807	47.203	21.5	0.0976	42.803	1.76	0.0032690	0.380	2.914	5.444 4
1996TK ₆₆	cClas	6.19 ± 0.12	28.62 ± 4.51	0.99 ± 0.06	0.68 ± 0.07	0.55 ± 0.10	1.28 ± 0.13	2.37 ± 0.15	1.67 ± 0.09	42.408	43.198	3.3	0.0092	42.803	0.27	0.0000792	0.058	3.085	5.848 3, 7, 11, 8	
1996TS ₆₆	hClas	5.94 ± 0.13	30.39 ± 6.07	1.03 ± 0.08	0.67 ± 0.11	0.64 ± 0.09	1.23 ± 0.14	2.22 ± 0.21	1.67 ± 0.16	38.325	50.061	7.3	0.1328	44.193	0.83	0.0006180	0.184	3.064	5.847 12, 7, 9	
1997CT ₂₉	cClas	6.28 ± 0.13	33.37 ± 6.57	—	—	—	—	—	—	—	45.070	1.0	0.0270	43.870	0.15	0.0000175	0.032	3.100	5.923 10, 8, 3	
1997CV ₂₉	hClas	7.16 ± 0.03	28.17 ± 2.06	1.21 ± 0.01	0.65 ± 0.02	—	—	—	—	—	44.396	8.0	0.0494	42.307	0.68	0.0004958	0.148	3.057	5.763 13	
1997QH ₄	hClas	6.99 ± 0.14	27.56 ± 5.79	1.09 ± 0.14	0.64 ± 0.09	0.63 ± 0.11	1.26 ± 0.14	2.41 ± 0.23	1.73 ± 0.17	39.281	46.454	13.2	0.0837	42.868	1.11	0.0013165	0.243	3.018	5.691 3, 7, 14, 5	
1997RT ₅	hClas	—	—	—	—	—	—	—	—	—	42.373	12.7	0.0238	41.387	1.03	0.0011825	0.222	3.015	5.626 5	
1997SZ ₁₀	2-1	8.09 ± 0.10	29.55 ± 6.02	1.14 ± 0.08	0.69 ± 0.10	0.57 ± 0.10	1.42 ± 0.16	—	—	—	65.819	11.8	0.3656	48.198	1.80	0.0025999	0.419	2.931	5.654 3, 15	
1998FS ₁₄₄	hClas	6.72 ± 0.10	19.80 ± 4.03	0.95 ± 0.04	0.59 ± 0.06	0.51 ± 0.09	1.22 ± 0.09	—	—	—	42.739	9.9	0.0227	41.789	0.80	0.0007123	0.173	3.042	5.707 13, 8, 10	
1998HL ₁₅₁	SDO	8.12 ± 0.15	15.96 ± 12.52	0.79 ± 0.35	0.42 ± 0.21	0.79 ± 0.21	1.17 ± 0.21	2.29 ± 0.33	1.19 ± 0.28	36.989	44.385	28.0	0.0909	40.687	2.32	0.0005523	0.479	2.784	5.044 6, 16	
1998KG ₂	cClas	6.12 ± 0.11	24.41 ± 5.40	1.04 ± 0.10	0.61 ± 0.10	0.61 ± 0.08	1.23 ± 0.11	2.21 ± 0.15	1.60 ± 0.16	41.025	45.647	0.8	0.0533	43.336	0.25	0.0000455	0.055	3.091	5.883 4, 5, 8, 17	
1998KS ₆₅	cClas	7.17 ± 0.04	27.07 ± 2.06	1.09 ± 0.04	0.64 ± 0.02	—	—	—	—	—	45.090	1.2	0.0319	43.697	0.17	0.0000243	0.038	3.097	5.911 13, 8	
1998UR ₄₃	Plut	8.08 ± 0.13	22.09 ± 7.92	0.78 ± 0.09	0.63 ± 0.12	0.48 ± 0.17	1.17 ± 0.17	—	—	—	48.775	8.8	0.2262	39.776	1.29	0.0013866	0.273	2.970	5.454 4, 14, 8	
1998UL ₁₃	4-3	6.98 ± 0.09	17.19 ± 5.40	0.95 ± 0.08	0.56 ± 0.09	0.50 ± 0.09	1.05 ± 0.09	2.02 ± 0.06	1.49 ± 0.05	31.780	41.455	9.6	0.1306	36.668	1.04	0.0010427	0.211	2.979	5.333 8, 18	
1998WS ₃₁	Plut	7.95 ± 0.19	18.40 ± 6.63	0.73 ± 0.08	0.57 ± 0.10	0.50 ± 0.14	1.10 ± 0.19	1.74 ± 0.10	1.31 ± 0.07	31.490	47.814	6.7	0.2058	39.652	1.12	0.0010151	0.237	2.990	5.497 8, 19	
1998WV ₂₄	cClas(b)	7.12 ± 0.07	12.00 ± 4.20	0.77 ± 0.01	0.50 ± 0.06	0.45 ± 0.11	0.96 ± 0.11	—	—	—	40.716	1.5	0.0389	39.193	0.27	0.0000420	0.047	3.048	5.616 3, 8	
1998WV ₃₁	Plut	7.62 ± 0.07	16.39 ± 5.55	0.79 ± 0.10	0.55 ± 0.08	0.48 ± 0.15	1.01 ± 0.17	1.81 ± 0.16	1.35 ± 0.10	28.767	50.156	5.7	0.2710	39.462	1.37	0.0014146	0.289	2.956	5.407 14, 19	
1998WX ₂₄	cClas	6.24 ± 0.10	30.43 ± 5.30	1.09 ± 0.05	0.73 ± 0.09	0.50 ± 0.09	1.29 ± 0.11	—	—	—	45.140	0.9	0.0363	43.559	0.18	0.0000249	0.040	3.096	5.902 3, 8	
1998WX ₃₁	cClas	6.17 ± 0.09	28.83 ± 5.13	—	—	—	—	—	—	—	50.931	3.0	0.1131	45.758	0.55	0.0002333	0.124	3.105	5.999 14, 8	
1998WY ₂₄	cClas	6.44 ± 0.09	27.89 ± 5.30	—	—	—	—	—	—	—	45.467	1.9	0.0457	43.475	0.26	0.0000556	0.057	3.093	5.891 8	
1998WZ ₃₁	Plut	8.04 ± 0.10	6.52 ± 6.87	0.73 ± 0.09	0.47 ± 0.12	0.35 ± 0.10	0.83 ± 0.14	1.59 ± 0.11	1.26 ± 0.09	32.928	46.635	14.6	0.1723	39.782	1.45	0.0020606	0.305	2.949	5.403 8, 19	
1998XJ ₆₅	SDTNO	6.43 ± 0.14	39.45 ± 8.35	0.93 ± 0.23	0.72 ± 0.14	0.75 ± 0.15	1.48 ± 0.15	2.35 ± 0.24	1.58 ± 0.21	37.297	92.522	6.7	0.4254	64.910	1.63	0.0019497	0.441	3.105	6.430 2, 8	
1999CB ₁₁₉	hClas	6.74 ± 0.05	34.30 ± 4.32	1.21 ± 0.10	0.71 ± 0.08	0.65 ± 0.06	1.36 ± 0.10	2.57 ± 0.10	1.93 ± 0.10	40.825	53.088	8.7	0.1306	46.957	0.87	0.0007194	0.200	3.089	5.998 19	
1999CD ₁₅₈	7-4	4.84 ± 0.11	16.73 ± 5.83	0.84 ± 0.07	0.53 ± 0.09	0.57 ± 0.11	1.10 ± 0.07	1.93 ± 0.09	1.38 ± 0.12	37.655	49.969	25.4	0.1405	43.812	2.09	0.0044889	0.452	2.845	5.308 14, 11, 16	
1999CF ₁₁₉	SDTNO	6.98 ± 0.08	18.78 ± 4.59	0.83 ± 0.06	0.62 ± 0.07	0.34 ± 0.11	1.00 ± 0.11	—	—	—	140.799	19.7	0.5691	89.732	2.09	0.0035201	0.661	3.010	6.488 14, 6, 8, 20	
1999CI ₁₁₉	cClas	6.70 ± 0.21	32.69 ± 26.17	1.38 ± 0.18	0.69 ± 0.28	—	—	—	—	—	48.750	3.2	0.0708	45.528	0.40	0.0001371	0.090	3.111	6.006 6	
1999CL ₁₁₉	hClas	5.62 ± 0.10	15.56 ± 4.51	—	—	—	—	—	—	—	47.492	23.3	0.0091	47.066	1.76	0.0033128	0.395	2.938	5.637 19	
1999CM ₁₁₉	cClas	7.36 ± 0.06	56.52 ± 23.48	0.90 ± 0.19	0.88 ± 0.26	—	—	—	—	—	47.362	2.7	0.0769	43.979	0.41	0.0001360	0.091	3.093	5.909 6	
1999CQ ₃₃	hClas	6.68 ± 0.05	22.75 ± 9.15	0.75 ± 0.07	0.60 ± 0.10	—	—	—	—	—	45.289	13.3	0.0929	41.438	1.15	0.0013989	0.247	3.001	5.595 6	
1999CX ₁₃₁	5-3	6.92 ± 0.07	19.49 ± 5.46	0.95 ± 0.09	0.60 ± 0.10	0.48 ± 0.07	1.07 ± 0.10	2.04 ± 0.08	1.54 ± 0.08	32.517	52.226	9.7	0.2326	42.371	1.32	0.0014739	0.288	2.985	5.593 8, 19, 18	
1999GS ₆₆	hClas	6.23 ± 0.02	25.97 ± 10.53	1.13 ± 0.09	0.63 ± 0.11	—	—	—	—	—	48.308	5.2	0.0863	44.469	0.56	0.0002882	0.125	3.089	5.918 6	
1999HG ₁₂	7-4	7.32 ± 0.06	25.21 ± 2.97	1.02 ± 0.08	0.59 ± 0.06	0.66 ± 0.03	1.25 ± 0.07	2.27 ± 0.06	1.61 ± 0.05	36.840	50.134	1.0	0.1528	43.487	0.69	0.0003430	0.154	3.068	5.833 18	
1999HI ₁₂	cClas	—	—	—	—	—	—	—	—	—	44.864	4.5	0.0456	42.908	0.42	0.0001766	0.091	3.080	5.840 8	
1999HS ₁₁	cClas	6.35 ± 0.08	33.64 ± 4.14	1.13 ± 0.08	0.72 ± 0.06	0.60 ± 0.09	1.28 ± 0.13	2.29 ± 0.21	1.84 ± 0.10	43.257	44.754	2.6	0.0170	44.005	0.22	0.0000509	0.048	3.100	5.928 17, 13, 19, 21	
1999HV ₁₁	cClas	7.01 ± 0.05	21.71 ± 2.06	1.11 ± 0.06	0.59 ± 0.02	—	—	—	—	—	43.982	3.2	0.0269	42.830	0.28	0.0000814	0.061	3.085	5.849 13	
1999ID ₁₃₂	hClas	5.99 ± 0.02	10.74 ± 12.41	1.11 ± 0.10	0.48 ± 0.13	—	—	—	—	—	46.106	10.5	0.0673	43.200	0.88	0.0008335	0.194	3.048	5.774 6	
1999OE ₄	cClas	6.89 ± 0.19	34.57 ± 12.26	0.88 ± 0.17	0.76 ± 0.20	0.55 ± 0.21	1.22 ± 0.32	2.46 ± 0.19	1.83 ± 0.15	43.224	47.406	2.2	0.0461	45.315	0.26	0.0006005	0.059	3.114	6.006 19, 8	
1999OH ₄	hClas	6.32 ± 0.12	—	—	—	—	—	—	—	—	41.919	28.2	0.0274	40.414	2.31	0.0005542	0.474	2.786	5.037 16	
1999OI ₄	SDO(b)	6.90 ± 0.06	27.37 ± 3.21	0.99 ± 0.08	0.68 ± 0.05	0.52 ± 0.06	1.22 ± 0.10	2.19 ± 0.10	1.68 ± 0.08	36.965	59.049	4.0	0.0274	38.007	0.36	0.0001406	0.075	3.033	5.527 19	
1999OM ₄	cClas	—	20.01 ± 5.97	1.14 ± 0.14	0.60 ± 0.10	0.50 ± 0.09	1.10 ± 0.14	2.24 ± 0.19	1.74 ± 0.17	40.511	51.338	2.1	0.11							

Table 5 cont'd

Object	Class ^a	H_R^b	Gr^c	$B-V$	$V-R$	$R-I$	$V-I$	$B-I$	$B-R$	q	Q	i	e	a	u_r^d	u^e	e^f	T_N^g	T_h	References
2000CF ₁₀₅	cClas(b)	6.47 ± 0.29	-0.68 ± 15.42	0.78 ± 0.09	0.33 ± 0.24	0.40 ± 0.30	0.52 ± 0.53	—	1.75 ± 0.12	42.232	45.876	0.5	0.0414	44.054	0.19	0.0000262	0.042	3.101	5.932	20, 49, 8
2000CG ₁₀₅	hClas	5.69 ± 0.07	33.69 ± 4.04	1.08 ± 0.08	0.68 ± 0.07	0.70 ± 0.06	1.38 ± 0.09	2.46 ± 0.12	1.17 ± 0.17	44.593	48.371	27.9	0.0406	46.482	2.14	0.0004769	0.470	2.842	5.388	6, 8, 16
2000CJ ₁₀₅	hClas	6.37 ± 0.12	24.31 ± 7.41	—	0.59 ± 0.13	0.65 ± 0.11	1.24 ± 0.11	—	1.76 ± 0.11	39.490	48.964	11.6	0.1071	44.227	1.02	0.0010739	0.228	3.042	5.797	15
2000CK ₁₀₅	Plut	6.40 ± 0.09	23.70 ± 6.24	1.22 ± 0.17	0.56 ± 0.11	0.68 ± 0.08	1.23 ± 0.11	2.56 ± 0.21	1.85 ± 0.19	40.985	48.576	8.1	0.2291	39.523	0.27	0.0013347	0.269	2.970	5.443	8
2000CL ₁₀₄	cClas	6.76 ± 0.06	4.33 ± 11.11	1.11 ± 0.08	0.41 ± 0.12	—	—	—	1.52 ± 0.09	41.302	45.007	4.2	0.0429	44.524	0.37	0.0000993	0.082	3.101	5.947	5, 8
2000CN ₁₀₅	cClas	5.29 ± 0.16	27.80 ± 5.69	1.06 ± 0.10	0.61 ± 0.10	0.69 ± 0.08	1.35 ± 0.11	2.41 ± 0.15	1.72 ± 0.13	40.262	49.010	3.4	0.0980	44.636	0.51	0.0002141	0.115	3.094	5.936	19, 15
2000CO ₁₀₅	hClas	5.62 ± 0.12	24.69 ± 6.88	0.82 ± 0.06	0.65 ± 0.13	0.55 ± 0.08	1.17 ± 0.08	—	1.52 ± 0.18	40.358	54.170	19.2	0.1461	47.264	1.58	0.0025769	0.360	2.978	5.740	6, 8
2000CP ₁₀₄	hClas	6.59 ± 0.06	19.27 ± 3.61	—	0.53 ± 0.06	0.64 ± 0.06	1.14 ± 0.10	—	1.72 ± 0.13	39.994	48.696	9.5	0.0981	44.345	0.86	0.0007470	0.192	3.062	5.848	15, 8
2000CQ ₁₀₄	4:3	7.84 ± 0.06	9.80 ± 3.19	0.83 ± 0.08	0.48 ± 0.06	0.44 ± 0.04	0.92 ± 0.07	1.75 ± 0.06	1.31 ± 0.05	28.091	45.438	13.5	0.2359	36.765	1.64	0.0024288	0.332	2.908	5.165	18
2000CQ ₁₀₅	SDO	6.00 ± 0.05	6.47 ± 4.15	0.68 ± 0.04	0.45 ± 0.06	0.35 ± 0.10	0.90 ± 0.14	1.53 ± 0.12	1.11 ± 0.04	34.737	79.544	19.7	0.3921	57.141	2.05	0.0036613	0.517	2.915	5.833	6, 8, 19, 20
2000CQ ₁₁₄	cClas(b)	6.66 ± 0.15	32.47 ± 8.98	—	0.69 ± 0.16	0.65 ± 0.13	1.38 ± 0.11	—	—	40.858	51.534	2.7	0.1156	46.196	0.55	0.0002284	0.125	3.111	6.025	8
2000FS ₅₃	cClas	7.17 ± 0.12	34.27 ± 7.12	1.09 ± 0.11	0.70 ± 0.08	—	1.31 ± 0.16	—	1.79 ± 0.10	41.505	44.422	2.1	0.0340	42.964	0.23	0.0000475	0.050	3.088	5.861	6, 13, 8
2000FU ₅₃	Plut	7.56 ± 0.07	13.94 ± 6.02	—	0.45 ± 0.09	0.68 ± 0.12	1.13 ± 0.15	—	—	32.751	45.422	17.4	0.1621	39.086	1.63	0.0026961	0.339	2.917	5.296	19
2000FZ ₅₃	Cent	11.11 ± 0.16	14.11 ± 5.09	0.61 ± 0.03	0.60 ± 0.08	0.29 ± 0.09	0.97 ± 0.12	—	1.17 ± 0.05	12.331	35.095	34.9	0.4800	23.713	4.73	0.0198494	0.746	2.547	3.293	19, 20
2000GV ₁₄₆	cClas	5.99 ± 0.10	23.51 ± 5.42	0.91 ± 0.04	0.58 ± 0.10	0.64 ± 0.06	1.08 ± 0.06	—	1.55 ± 0.05	37.558	44.701	19.1	0.0868	41.130	1.60	0.0027273	0.339	2.933	5.419	13, 8
2000KK ₁	hClas	—	—	—	—	—	—	—	—	36.036	40.901	21.3	0.0632	38.468	1.81	0.0034997	0.369	2.885	5.191	8
2000KL ₄	SDO	6.80 ± 0.16	35.05 ± 10.05	—	0.61 ± 0.16	0.92 ± 0.19	1.62 ± 0.16	—	—	41.088	45.102	4.4	0.0466	43.095	0.41	0.0001694	0.090	3.082	5.853	19, 22
2000QU ₆₉	cClas	—	—	—	—	—	1.13 ± 0.15	—	—	45.511	47.435	4.6	0.0207	46.473	0.36	0.0001444	0.083	3.125	5.854	17, 23, 24
2000PD ₃₀	SDO	5.87 ± 0.11	2.78 ± 4.11	0.75 ± 0.07	0.37 ± 0.08	0.41 ± 0.05	0.76 ± 0.09	1.56 ± 0.06	1.13 ± 0.08	35.766	72.618	18.4	0.3400	54.192	1.89	0.0031792	0.464	2.950	5.854	17, 23, 24
2000PE ₃₀	cClas	—	—	—	—	—	—	—	—	41.880	46.414	3.7	0.0514	44.147	0.31	0.0000862	0.069	3.099	5.929	21
2000QC ₂₂₆	cClas	—	45.97 ± 3.79	1.08 ± 0.05	0.80 ± 0.04	—	—	—	1.88 ± 0.06	41.880	58.240	2.7	0.0216	47.873	0.97	0.0006978	0.226	3.087	6.019	18
2000QJ ₂₅₁	2:1(b)	6.59 ± 0.04	11.64 ± 2.46	0.87 ± 0.06	0.49 ± 0.04	0.48 ± 0.03	0.93 ± 0.09	1.84 ± 0.05	1.36 ± 0.04	37.505	58.240	3.7	0.1257	42.302	0.58	0.0002342	0.126	3.064	5.780	18
2000QN ₂₅₁	5:3	7.66 ± 0.09	22.96 ± 5.23	1.04 ± 0.12	0.59 ± 0.09	0.61 ± 0.07	1.20 ± 0.12	2.24 ± 0.10	1.63 ± 0.07	36.986	47.618	0.3	0.1257	42.302	0.58	0.0002342	0.126	3.064	5.780	18
2000YB ₂₉	SDO	6.43 ± 0.08	24.54 ± 5.05	0.86 ± 0.10	0.64 ± 0.09	0.55 ± 0.07	1.19 ± 0.11	2.05 ± 0.15	1.50 ± 0.13	27.076	52.073	7.7	0.3158	39.575	1.62	0.0020729	0.343	2.917	5.318	15
2001FL ₁₈₅	cClas(b)	—	—	—	—	—	1.19 ± 0.23	—	—	40.770	47.602	3.6	0.0770	44.186	0.44	0.0001718	0.099	3.093	5.917	49
2001FM ₉₄	SDO	7.46 ± 0.16	12.31 ± 2.98	0.76 ± 0.03	0.45 ± 0.04	0.62 ± 0.06	1.09 ± 0.08	—	1.19 ± 0.04	34.316	73.625	28.7	0.3642	53.970	2.51	0.0058073	0.603	2.746	5.359	20, 15
2001FQ ₁₈₅	2:1	7.43 ± 0.06	35.25 ± 3.19	1.18 ± 0.08	0.68 ± 0.06	0.73 ± 0.04	1.41 ± 0.07	2.59 ± 0.06	1.86 ± 0.05	36.749	58.464	3.2	0.2281	47.606	1.01	0.0007500	0.235	3.078	5.990	18
2001FU ₁₇₂	Plut	8.17 ± 0.18	24.96 ± 12.06	—	0.67 ± 0.19	0.47 ± 0.23	1.14 ± 0.24	—	—	28.731	49.849	24.8	0.2867	39.290	2.41	0.0056125	0.498	2.765	4.939	16
2001HY ₆₅	hClas	6.04 ± 0.06	18.33 ± 4.03	1.03 ± 0.06	0.48 ± 0.07	0.71 ± 0.06	1.19 ± 0.09	2.22 ± 0.11	1.51 ± 0.09	37.917	48.389	17.1	0.1213	43.153	1.46	0.0022271	0.319	2.969	5.584	15
2001HZ ₅₈	cClas	6.16 ± 0.05	16.87 ± 3.64	1.17 ± 0.06	0.47 ± 0.06	0.70 ± 0.06	1.17 ± 0.09	2.34 ± 0.10	1.64 ± 0.09	40.890	44.331	2.9	0.0404	42.610	0.30	0.0000857	0.065	3.081	5.833	15
2001KA ₇₇	hClas	5.05 ± 0.09	36.04 ± 4.69	1.10 ± 0.11	0.71 ± 0.07	0.72 ± 0.12	1.45 ± 0.09	2.55 ± 0.16	1.81 ± 0.12	42.797	51.685	11.9	0.0941	47.241	0.99	0.0010237	0.227	3.078	5.980	11, 19, 22
2001KB ₇₇	Plut	7.35 ± 0.08	12.71 ± 4.52	0.89 ± 0.10	0.47 ± 0.08	0.56 ± 0.07	1.01 ± 0.10	—	1.39 ± 0.13	28.390	50.467	17.5	0.2800	39.428	1.97	0.0035464	0.411	2.859	5.171	19, 25
2001KD ₇₇	Plut	5.93 ± 0.10	23.50 ± 5.13	1.12 ± 0.05	0.62 ± 0.07	0.56 ± 0.07	1.20 ± 0.07	2.31 ± 0.09	1.76 ± 0.06	35.005	43.787	2.3	0.1115	39.396	0.56	0.0002363	0.118	3.036	5.997	11, 19, 23
2001KG ₇₆	9:4	—	45.97 ± 5.60	1.12 ± 0.06	0.80 ± 0.06	—	—	—	1.92 ± 0.09	33.846	68.902	1.6	0.3412	51.374	1.42	0.0014304	0.342	3.042	6.006	21
2001KG ₇₇	SDO	8.34 ± 0.12	7.03 ± 5.59	0.81 ± 0.04	0.44 ± 0.06	—	—	—	1.24 ± 0.07	33.950	88.690	15.5	0.4464	61.320	1.98	0.0031961	0.520	2.953	6.005	20
2001KY ₇₆	Plut	6.69 ± 0.18	7.47 ± 8.54	1.45 ± 0.22	0.41 ± 0.13	0.54 ± 0.18	1.03 ± 0.15	2.51 ± 0.24	1.96 ± 0.29	29.973	48.446	4.0	0.2356	39.209	1.47	0.0010071	0.246	2.981	5.455	22
2001QJ ₂₉₇	hClas(b)	6.38 ± 0.17	7.90 ± 4.21	0.66 ± 0.07	0.37 ± 0.07	0.63 ± 0.07	0.85 ± 0.12	1.66 ± 0.12	1.03 ± 0.10	40.582	52.086	30.6	0.1241	46.334	2.40	0.0057986	0.524	2.769	5.210	15
2001QJ ₂₉₈	hClas	6.18 ± 0.17	30.40 ± 8.33	0.97 ± 0.13	0.67 ± 0.09	—	—	—	1.64 ± 0.16	40.280	44.921	5.0	0.0545	42.601	0.47	0.0002236	0.103	3.074	5.814	22
2001QJ ₂₉₈	7:4	7.39 ± 0.10	31.11 ± 5.35	1.20 ± 0.13	0.69 ± 0.10	0.62 ± 0.06	1.31 ± 0.12	2.51 ± 0.10	1.89 ± 0.08	36.917	50.766	3.7	0.1579	43.841	1.97	0.0004485	0.170	3.066	5.839	18
2001QJ ₂₉₈	Plut	5.12 ± 0.12	4.60 ± 4.64	0.67 ± 0.07	0.39 ± 0.06	0.57 ± 0.18	0.89 ± 0.19	1.34 ± 0.09	1.05 ± 0.09	35.118	43.699	22.4	0.1089	39.408	1.92	0.0003869	0.396	2.868	5.192	26, 23, 22,
2001QJ ₂₉₈	25	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
2001QQ ₂₉₇	cClas	5.97 ± 0.10	29.79 ± 6.44	—	0.69 ± 0.11	0.58 ± 0.10	1.27 ± 0.15	—	—	41.302	44.681	1.1	0.0393	42.991	0.20	0.0000316	0.044	3.089	5.864	15
2001QP ₂₉₇	cClas	6.68 ± 0.09	25.64 ± 5.83	—	0.63 ± 0.10	0.60 ± 0.09	1.23 ± 0.13	—	—	39.993	50.554	1.4	0.1166	45.273	0.53	0.0002016	0.119	3.101	5.972	15
2001QX ₃₂₂	SDO	6.14 ± 0.15	19.06 ± 5.95	1.13 ± 0.24	0.62 ± 0.10	0.42 ± 0.09	1.07 ± 0.14	2.43 ± 0.19	1.75 ± 0.28	35.668	80.930	28.5	0.3882	58.299	2.46	0.0055191	0.615	2.771	5.511	20, 15
2001QX ₃₂₂	cClas(b)	5.15 ± 0.23	24.41 ± 8.17	0.94 ± 0.13	0.61 ± 0.10	0.65 ± 0.29	1.34 ± 0.20	1.78 ± 0.34	1.56 ± 0.18	40.210	47.302	1.5	0.0810	43.756	0.35	0.0001105	0.085	3.099	5.937	22, 23, 21
2001RZ ₁₄₃	cClas(b)	6.24 ± 0.12	13.26 ± 7.24	1.08 ± 0.14	0.51 ± 0.13	0.49 ± 0.10	1.00 ± 0.16	2.08 ± 0.21	1.59 ± 0.10	41.255	47.272	2.1	0.0680	44.264	0.38	0.0000963	0.077	3.099	5.893	15
2001UP ₁₈	2:1	5.96 ± 0.07	16.55 ± 3.94	0.95 ± 0.09	0.54 ± 0.07	0.53 ± 0.05	1.07 ± 0.09	2.02 ± 0.07	1.49 ± 0.05	44.504	51.324	1.2	0.0712	47.914	0.32	0.0000748	0.074	3.145	6.161	18
2001XP ₂₅₄	5:3(b)	7.54 ± 0.04	24.86 ± 2.66	1.09 ± 0.06	0.62 ± 0.04	0.60 ± 0.04	1.22 ± 0.06	2.31 ± 0.06	1.71 ± 0.04	33.043	51.979	2.6	0.2227	45.961	1.84	0.0007778	0.227	3.023	5.690	18
2001XQ ₂₅₄	5:2	7.67 ± 0.05	10.84 ± 2.6																	

Table 5 cont'd

Object	Class ^a	H_R^b	GrF^c	$B-V$	$V-R$	$R-I$	$V-I$	$B-I$	$B-R$	q	Q	i	e	a	u_v^d	u_r^e	e^f	T_N^g	T_h^h	References	
2002VT ₁₃₀	cClas(b)	5.42 ± 0.09	21.30 ± 5.83	1.45 ± 0.21	0.56 ± 0.10	0.63 ± 0.09	1.19 ± 0.13	2.64 ± 0.25	2.01 ± 0.23	41.398	43.983	1.2	0.0303	42.691	0.17	0.0000230	0.036	3.086	5.847	15	
2002VV ₁₃₀	5:3	7.55 ± 0.03	34.21 ± 1.92	1.22 ± 0.05	0.70 ± 0.04	0.67 ± 0.02	1.37 ± 0.04	2.59 ± 0.04	1.92 ± 0.03	35.156	50.013	2.4	0.1744	42.585	0.82	0.0004871	0.179	3.048	5.751	18	
2002XX ₆₃	Plut	4.43 ± 0.04	0.85 ± 2.06	0.72 ± 0.02	0.37 ± 0.02	—	—	—	—	0.09 ± 0.03	34.623	44.477	13.3	0.1246	39.550	1.24	0.0015768	0.261	2.975	5.457	25
2003AZ ₆₄	Plut(b)	3.54 ± 0.05	3.65 ± 3.46	0.65 ± 0.04	0.38 ± 0.04	0.55 ± 0.14	0.83 ± 0.12	1.50 ± 0.18	1.05 ± 0.06	32.730	46.552	13.5	0.1743	39.641	1.39	0.0018618	0.292	2.957	5.416	26, 15, 25, 27, 28	
2003FF ₁₂₈	Plut	—	34.99 ± 2.91	1.05 ± 0.04	0.71 ± 0.03	—	—	—	1.76 ± 0.05	31.022	47.729	1.9	0.2121	39.376	1.02	0.0007426	0.215	2.999	5.506	21	
2003FZ ₁₂₉	SDTNO	6.98 ± 0.04	10.40 ± 2.26	0.84 ± 0.06	0.48 ± 0.04	0.46 ± 0.03	0.94 ± 0.05	1.78 ± 0.05	1.32 ± 0.04	37.986	85.415	5.8	0.3843	61.700	1.51	0.0016617	0.397	3.119	6.410	24	
2003HB ₅₇	SDTNO	7.39 ± 0.03	12.70 ± 1.89	0.83 ± 0.04	0.48 ± 0.03	0.54 ± 0.03	1.02 ± 0.04	1.85 ± 0.04	1.31 ± 0.03	38.088	279.554	15.5	0.7602	158.821	1.90	0.0027244	0.806	3.067	6.950	24	
2003HX ₅₆	SDO	7.03 ± 0.21	—	1.58 ± 16.51	—	0.35 ± 0.23	0.26 ± 0.46	0.61 ± 0.46	—	36.504	57.906	29.6	0.2267	47.205	2.44	0.0058340	0.543	2.760	5.214	16	
2003QA ₉₂	SDO	6.37 ± 0.24	25.97 ± 3.79	1.04 ± 0.03	0.63 ± 0.04	—	—	—	1.67 ± 0.02	35.879	40.556	3.4	0.0612	38.218	0.41	0.0001550	0.086	3.033	5.537	29	
2003QK ₆₁	SDTNO	6.97 ± 0.04	11.97 ± 2.27	0.87 ± 0.06	0.50 ± 0.04	0.47 ± 0.03	0.97 ± 0.05	1.84 ± 0.05	1.37 ± 0.04	38.447	96.796	4.0	0.4314	67.621	1.58	0.0017929	0.437	3.144	6.565	24	
2003QO ₉₁	SDO	7.62 ± 0.28	13.62 ± 7.42	0.67 ± 0.06	0.51 ± 0.08	—	—	—	1.18 ± 0.05	36.006	41.753	5.4	0.0739	38.879	0.57	0.0003156	0.120	3.031	5.562	29	
2003QW ₁₁₁	7:4(b)	6.74 ± 0.07	24.54 ± 3.88	1.07 ± 0.10	0.61 ± 0.08	0.61 ± 0.04	1.25 ± 0.06	2.29 ± 0.08	1.68 ± 0.07	38.739	48.766	2.7	0.1146	43.752	0.56	0.0002370	0.124	3.081	5.814	18	
2003QW ₉₀	hClas	4.73 ± 0.06	30.80 ± 3.43	1.11 ± 0.07	0.67 ± 0.06	0.65 ± 0.05	1.32 ± 0.08	2.43 ± 0.10	1.78 ± 0.09	40.441	47.338	10.4	0.0786	43.890	0.88	0.0008239	0.196	3.055	5.875	15	
2003QY ₁₁₁	cClas	7.27 ± 0.16	17.53 ± 7.73	—	—	0.46 ± 0.16	0.69 ± 0.07	1.15 ± 0.17	—	41.833	44.980	2.9	0.0362	43.406	0.28	0.0000799	0.063	3.091	5.885	15	
2003SQ ₃₁₇	hClas(h)	6.14 ± 0.02	—	—	—	—	—	—	—	39.238	46.256	28.6	0.0821	42.747	2.30	0.0054474	0.485	2.791	5.139	16	
2003TH ₁₆₈	Plut	6.94 ± 0.06	4.60 ± 3.88	0.61 ± 0.06	0.38 ± 0.06	0.48 ± 0.06	0.86 ± 0.06	1.47 ± 0.06	0.99 ± 0.07	35.805	43.389	27.9	0.0958	39.597	2.34	0.0056668	0.477	2.779	4.986	16	
2003TL ₅₈	cClas(b)	—	—	—	—	—	1.13 ± 0.04	—	—	40.596	48.820	1.0	0.0920	44.708	0.42	0.0001245	0.093	3.101	5.953	49	
2003UY ₂₉₁	dTNO	—	18.76 ± 4.21	0.88 ± 0.10	0.51 ± 0.07	0.67 ± 0.07	1.18 ± 0.10	2.06 ± 0.10	1.39 ± 0.07	41.360	57.790	3.5	0.1657	49.575	0.75	0.0004232	0.177	3.134	6.181	24	
2003UZ ₁₁₇	hClas(h)	4.92 ± 0.08	0.16 ± 2.34	0.63 ± 0.05	0.36 ± 0.03	0.34 ± 0.06	0.70 ± 0.05	—	0.99 ± 0.05	38.362	50.655	27.4	0.1381	44.509	2.22	0.0050248	0.480	2.815	5.261	25, 27	
2003UZ ₁₁₃	SDO	4.03 ± 0.04	6.64 ± 3.72	—	0.45 ± 0.06	0.39 ± 0.07	0.82 ± 0.06	—	—	30.731	48.032	12.0	0.2197	39.381	1.44	0.0018712	0.303	2.947	5.382	27	
2003YL ₁₇₉	cClasi	7.48 ± 0.30	7.95 ± 9.24	0.81 ± 0.04	0.45 ± 0.10	—	—	—	1.26 ± 0.09	38.740	39.357	2.5	0.0079	39.048	0.21	0.0000503	0.045	3.047	5.607	29	
2003YW ₁₇₉	5:3	7.13 ± 0.06	32.37 ± 3.19	1.23 ± 0.08	0.71 ± 0.06	0.61 ± 0.04	1.32 ± 0.07	2.55 ± 0.06	1.94 ± 0.05	35.694	49.144	2.4	0.1585	42.419	0.75	0.0004111	0.164	3.052	5.756	18	
2004EG ₉₆	4:3	7.91 ± 0.07	12.49 ± 3.48	0.88 ± 0.09	0.51 ± 0.07	0.47 ± 0.03	0.98 ± 0.08	1.86 ± 0.07	1.39 ± 0.06	32.053	78.700	16.2	0.4212	55.377	2.03	0.0034117	0.505	2.907	5.776	18	
2004HM ₇₉	4:3	7.38 ± 0.04	10.97 ± 4.48	0.82 ± 0.11	0.47 ± 0.08	0.50 ± 0.06	0.97 ± 0.10	1.79 ± 0.09	1.29 ± 0.07	33.407	39.362	1.2	0.0818	36.384	0.42	0.0001266	0.084	3.019	5.413	18	
2004HO ₇₉	5:2	7.10 ± 0.06	22.64 ± 5.98	0.97 ± 0.14	0.55 ± 0.11	0.68 ± 0.08	1.23 ± 0.12	2.20 ± 0.12	1.52 ± 0.09	32.397	77.666	5.6	0.4413	55.303	1.70	0.0020962	0.423	3.001	5.995	18	
2004OI ₁₄	SDTNO	6.99 ± 0.03	15.45 ± 1.89	0.90 ± 0.04	0.52 ± 0.03	0.54 ± 0.03	1.06 ± 0.04	1.96 ± 0.04	1.42 ± 0.03	39.211	70.983	22.5	0.2883	55.097	1.95	0.0036026	0.479	2.941	5.851	24	
2004OQ ₁₅	7:4	6.99 ± 0.04	9.54 ± 3.94	0.87 ± 0.09	0.49 ± 0.07	0.41 ± 0.05	0.90 ± 0.09	1.77 ± 0.08	1.36 ± 0.06	38.141	48.934	9.7	0.1240	43.537	0.95	0.0008784	0.210	3.044	5.777	18	
2004PB ₁₀₈	hClas(b)	—	—	—	—	—	1.12 ± 0.07	—	—	40.136	50.184	20.2	0.1113	45.160	1.64	0.0028163	0.363	2.952	5.610	49	
2004PT ₁₀₇	hClas	5.84 ± 0.03	—	—	—	—	—	—	—	38.187	42.877	26.2	0.0579	40.532	2.15	0.0048470	0.445	2.823	5.130	16	
2004PW ₁₀₇	7:4	7.61 ± 0.18	25.18 ± 10.42	1.15 ± 0.24	0.66 ± 0.19	0.53 ± 0.13	1.19 ± 0.23	2.34 ± 0.19	1.81 ± 0.14	37.942	49.721	2.8	0.1344	43.832	0.64	0.0003126	0.143	3.076	5.864	18	
2004TX ₃₅₇	SDO	7.61 ± 0.06	15.50 ± 3.41	0.95 ± 0.07	0.55 ± 0.06	0.48 ± 0.04	1.03 ± 0.07	1.98 ± 0.06	1.50 ± 0.04	31.494	80.391	9.0	0.4370	55.943	1.85	0.0025680	0.464	2.961	5.919	18	
2004TX ₃₅₇	2:1	6.53 ± 0.03	1.80 ± 1.92	0.68 ± 0.04	0.39 ± 0.04	0.34 ± 0.02	0.73 ± 0.04	1.41 ± 0.03	1.07 ± 0.02	34.500	61.748	9.7	0.2831	48.124	1.42	0.0016352	0.330	3.017	5.858	18	
2004TX ₃₅₇	4:3	8.54 ± 0.07	15.94 ± 3.71	0.99 ± 0.09	0.56 ± 0.08	0.48 ± 0.03	1.04 ± 0.08	2.03 ± 0.06	1.55 ± 0.05	28.869	44.837	16.2	0.2166	36.853	1.75	0.0029189	0.354	2.978	5.130	18	
2004VN ₁₁₂	dTNO	—	12.87 ± 3.63	0.90 ± 0.08	0.52 ± 0.06	0.45 ± 0.06	0.97 ± 0.08	1.87 ± 0.09	1.42 ± 0.06	47.331	63.905	25.5	0.8621	343.268	1.56	0.0018936	0.964	3.181	7.444	24	
2004XR ₉₀	dTNO	3.94 ± 0.04	10.10 ± 2.46	0.79 ± 0.06	0.45 ± 0.04	0.52 ± 0.04	0.97 ± 0.06	1.76 ± 0.06	1.24 ± 0.04	51.561	63.996	46.5	0.1076	57.778	3.21	0.0092397	0.734	2.417	4.649	24	
2005CA ₉₉	SDTNO	4.88 ± 0.04	23.62 ± 2.46	1.12 ± 0.05	0.64 ± 0.04	0.53 ± 0.03	1.17 ± 0.05	2.29 ± 0.04	1.76 ± 0.04	37.093	59.070	11.7	0.2285	48.081	1.31	0.0015294	0.305	3.037	5.905	18	
2005CB ₇₉	hClas(h)	4.37 ± 0.03	0.53 ± 2.25	0.74 ± 0.04	0.34 ± 0.04	0.38 ± 0.04	0.74 ± 0.04	1.47 ± 0.04	1.09 ± 0.03	37.281	49.679	28.6	0.1426	43.408	2.34	0.0055593	0.499	2.782	5.144	16	
2005EO ₂₉₇	3:1	7.22 ± 0.05	12.91 ± 3.49	0.84 ± 0.07	0.48 ± 0.05	0.57 ± 0.08	1.05 ± 0.09	1.89 ± 0.09	1.32 ± 0.05	41.121	85.006	25.0	0.3479	63.063	2.09	0.0040352	0.548	2.938	5.998	24	
2005EO ₃₀₄	cClas(b)	—	—	—	—	—	1.43 ± 0.14	—	—	42.578	48.589	3.4	0.0659	45.583	0.39	0.0001375	0.089	3.112	6.010	49	
2005ER ₃₁₈	SDO	7.68 ± 0.05	11.02 ± 2.66	0.84 ± 0.06	0.48 ± 0.05	0.48 ± 0.03	0.96 ± 0.06	1.80 ± 0.05	1.32 ± 0.04	30.636	42.586	10.4	0.1632	36.611	1.20	0.0013466	0.243	2.963	5.290	18	
2005GE ₁₈₇	Plut	7.19 ± 0.10	35.29 ± 6.49	0.98 ± 0.13	0.76 ± 0.10	0.55 ± 0.12	1.31 ± 0.14	2.29 ± 0.15	1.74 ± 0.11	26.557	51.845	18.3	0.3225	39.201	2.15	0.0041650	0.450	2.820	5.067	16	
2005PU ₂₁	SDO	6.09 ± 0.02	30.44 ± 1.43	1.14 ± 0.03	0.65 ± 0.02	0.68 ± 0.02	1.33 ± 0.03	2.47 ± 0.03	1.79 ± 0.02	29.225	327.810	6.2	0.8363	178.518	1.88	0.0025133	0.843	2.825	6.415	24	
2005RS ₁₃	hClas	4.92 ± 0.04	11.95 ± 2.46	0.87 ± 0.06	0.50 ± 0.04	0.47 ± 0.03	0.97 ± 0.05	1.84 ± 0.05	1.37 ± 0.04	38.450	57.461	10.0	0.1982	47.955	1.13	0.0011401	0.263	3.065	5.969	18	
2005SC ₂₇₈	SDO	6.70 ± 0.06	37.04 ± 2.97	1.29 ± 0.08	0.74 ± 0.06	0.66 ± 0.03	1.40 ± 0.07	2.69 ± 0.06	2.03 ± 0.05	32.531	37.333	1.5	0.0687	34.932	0.37	0.0001046	0.074	3.011	5.317	18	
2005SD ₂₇₈	SDTNO	5.91 ± 0.02	17.94 ± 1.43	0.97 ± 0.03	0.56 ± 0.02	0.53 ± 0.02	1.09 ± 0.03	2.06 ± 0.03	1.52 ± 0.02	39.946	71.884	17.9	0.2885	55.915	1.68	0.0025924	0.419	3.026	6.074	24	
2005SE ₂₇₈	hClas	6.76 ± 0.07	32.35 ± 3.72	1.16 ± 0.09	0.66 ± 0.07	0.70 ± 0.04	1.36 ± 0.08	2.52 ± 0.06	1.82 ± 0.05	37.685	47.295	6.9	0.1131	42.490	0.75	0.0005265	0.165	3.053	5.760	18	
2005SF ₂₇₈	SDO	6.27 ± 0.06	35.37 ± 3.19	1.27 ± 0.08	0.73 ± 0.06	0.64 ± 0.04	1.37 ± 0.07	2.64 ± 0.06	2.00 ± 0.05	35.525	52.484	13.3	0.1927	44.004	1.35	0.0017348	0.300	2.994	5.672	18	
2005UQ ₁₃	hClas	3.45 ± 0.15	—	—	—	—	—	—	—	37.151	49.778	25.7	0.1453	43.464	2.13</						

Table 5 cont'd

Object	Class ^a	H_R^b	Gr^c	$B-V$	$V-R$	$R-I$	$V-I$	$B-I$	$B-R$	q	Q	i	e	a	u_d	u^d	e^f	T_N^g	T_h^h	References
2007VL ₅₀₅	1:1	8.17 ± 0.05	11.78 ± 5.35	0.79 ± 0.13	0.45 ± 0.10	0.56 ± 0.06	1.01 ± 0.12	1.80 ± 0.10	1.24 ± 0.08	28.130	32.220	28.1	0.0678	30.175	2.68	0.0074503	0.476	2.760	4.411	18
2008FC ₇₆	Cent	9.18 ± 0.04	33.66 ± 3.86	—	0.71 ± 0.06	0.63 ± 0.08	1.34 ± 0.07	—	1.76 ± 0.02	10.175	19.350	27.1	0.3107	14.763	4.38	0.0181502	0.552	3.223	3.203	27
2008KV ₁₂	SDO(r)	8.56 ± 0.06	8.75 ± 3.63	0.82 ± 0.08	0.47 ± 0.06	0.42 ± 0.06	0.89 ± 0.08	1.71 ± 0.09	1.29 ± 0.06	21.159	61.993	103.5	0.4911	41.576	8.64	0.0263647	1.089	0.245	-1.025	24
2008OG ₁₉	SDTNO	4.61 ± 0.01	18.30 ± 0.94	0.94 ± 0.02	0.53 ± 0.01	0.59 ± 0.01	1.12 ± 0.02	2.06 ± 0.01	1.47 ± 0.01	38.577	93.869	13.1	0.4175	66.223	1.74	0.0024263	0.475	3.080	6.392	24
2008SJ ₂₃₆	Cent	12.39 ± 0.04	21.77 ± 2.74	—	0.62 ± 0.04	0.50 ± 0.05	1.12 ± 0.05	—	—	6.167	15.834	6.0	0.4394	11.000	4.05	0.0119753	0.452	3.814	3.071	27
2008YB ₃	Cent(r)	9.71 ± 0.01	10.24 ± 0.94	0.80 ± 0.02	0.46 ± 0.01	0.49 ± 0.01	0.95 ± 0.02	1.75 ± 0.01	1.26 ± 0.01	6.490	16.791	105.0	0.4425	11.640	16.44	0.0906494	1.062	2.294	-0.248	24
(2060) Chiron	Cent	6.31 ± 0.02	2.10 ± 1.28	0.64 ± 0.04	0.40 ± 0.01	0.31 ± 0.03	0.66 ± 0.03	1.28 ± 0.08	1.01 ± 0.04	8.640	18.846	6.9	0.3804	13.653	3.21	0.0076890	0.399	3.440	3.355	1, 30, 31, 32
(5145) Pholus	Cent	7.22 ± 0.06	47.65 ± 4.64	1.19 ± 0.10	0.78 ± 0.07	0.81 ± 0.09	1.52 ± 0.14	—	1.97 ± 0.11	8.467	18.447	24.7	0.5740	20.297	4.74	0.0187643	0.710	2.704	3.194	1, 33, 31, 27
(7066) Nessus	Cent	9.04 ± 0.07	42.41 ± 4.30	1.05 ± 0.11	0.78 ± 0.07	0.69 ± 0.08	1.43 ± 0.11	2.40 ± 0.21	1.85 ± 0.17	11.797	37.170	15.6	0.5182	24.484	3.52	0.0098236	0.584	2.714	3.786	1, 33, 31
(8405) Asbolus	Cent	9.26 ± 0.12	12.55 ± 2.12	0.70 ± 0.08	0.50 ± 0.03	0.50 ± 0.04	0.95 ± 0.06	1.64 ± 0.08	1.23 ± 0.06	6.880	29.257	17.6	0.6192	18.068	4.84	0.0183355	0.689	2.824	3.077	33, 31, 34, 35
(10199) Chariklo	Cent	6.58 ± 0.01	13.14 ± 1.30	0.81 ± 0.05	0.49 ± 0.02	0.52 ± 0.01	1.02 ± 0.02	1.86 ± 0.05	1.30 ± 0.07	13.051	18.424	23.4	0.1707	15.738	3.32	0.0111850	0.432	3.219	3.476	7, 36, 31
(10370) Hylonome	Cent	9.26 ± 0.13	10.93 ± 9.39	0.69 ± 0.07	0.47 ± 0.14	0.50 ± 0.19	0.92 ± 0.18	1.68 ± 0.13	1.15 ± 0.08	18.828	31.091	4.1	0.2457	24.959	1.52	0.0017206	0.256	2.966	4.443	1, 9, 31, 11, 9
(15760) 1992QB ₁	cClas	6.86 ± 0.12	37.35 ± 7.09	0.95 ± 0.13	0.73 ± 0.10	0.67 ± 0.20	1.24 ± 0.16	2.28 ± 0.27	1.67 ± 0.14	40.933	47.293	2.2	0.0721	44.113	0.37	0.0001067	0.082	3.096	5.922	1, 7, 3, 2, 8
(15788) 1993SB	Plut	8.03 ± 0.12	11.10 ± 5.75	0.80 ± 0.07	0.47 ± 0.08	0.52 ± 0.14	1.01 ± 0.14	1.77 ± 0.17	1.28 ± 0.10	26.745	52.519	1.9	0.3252	39.632	1.54	0.0016962	0.327	2.929	5.348	4, 7, 3, 14
(15789) 1993SC	Plut	6.71 ± 0.07	36.84 ± 4.88	1.04 ± 0.13	0.70 ± 0.08	0.73 ± 0.07	1.45 ± 0.09	2.38 ± 0.14	1.72 ± 0.14	32.333	47.009	5.2	0.1850	39.671	0.97	0.0007429	0.206	3.006	5.536	1, 37, 12, 7, 8
(15807) 1994GV ₉	cClas	6.82 ± 0.09	—	—	0.74 ± 0.10	—	—	—	—	40.992	46.469	0.6	0.0626	43.731	0.29	0.0000582	0.063	3.095	5.905	4
(15809) 1994JS	5:3	7.29 ± 0.10	39.25 ± 7.52	—	0.82 ± 0.12	0.48 ± 0.15	1.24 ± 0.16	—	—	33.043	51.025	14.1	0.2139	42.034	1.49	0.0020848	0.324	2.956	5.510	1, 8
(15810) 1994JR ₁	Plut	6.87 ± 0.08	15.17 ± 7.52	1.01 ± 0.18	0.52 ± 0.13	0.54 ± 0.12	1.01 ± 0.12	2.13 ± 0.22	1.61 ± 0.22	34.739	43.698	3.8	0.1142	39.219	0.63	0.0003201	0.132	3.031	5.575	9, 38, 9
(15820) 1994TB	Plut	7.52 ± 0.09	35.86 ± 6.85	1.09 ± 0.14	0.68 ± 0.12	0.75 ± 0.11	1.45 ± 0.13	2.55 ± 0.16	1.76 ± 0.15	26.966	52.544	12.1	0.3217	39.755	1.82	0.0027391	0.384	2.885	5.248	1, 37, 38, 7, 14
(15836) 1995DA ₂	4:3	7.99 ± 0.12	15.87 ± 10.88	—	0.54 ± 0.17	0.51 ± 0.21	1.05 ± 0.18	—	—	33.671	39.327	6.5	0.0775	36.499	0.68	0.0004592	0.138	3.006	5.389	9, 7
(15874) 1996TL ₄₆	SDO	5.12 ± 0.14	3.75 ± 3.46	0.71 ± 0.06	0.39 ± 0.06	0.35 ± 0.07	0.76 ± 0.09	1.34 ± 0.10	1.11 ± 0.07	35.053	134.413	24.0	0.5863	84.733	2.33	0.0044809	0.713	2.840	6.036	38, 12, 2, 7, 9
(15875) 1996TP ₆₆	Plut	6.94 ± 0.07	30.36 ± 4.74	1.03 ± 0.11	0.66 ± 0.07	0.66 ± 0.10	1.29 ± 0.11	2.25 ± 0.15	1.68 ± 0.12	26.396	53.028	5.7	0.3353	39.712	1.65	0.0020168	0.350	2.912	5.311	12, 2, 7, 38, 24, 9
(15883) 1997CR ₂₉	hClas	7.08 ± 0.14	19.00 ± 9.84	0.79 ± 0.10	0.54 ± 0.16	0.62 ± 0.18	1.17 ± 0.20	1.94 ± 0.18	1.26 ± 0.13	37.069	57.296	19.1	0.2143	47.183	1.72	0.0028841	0.392	2.949	5.668	7, 17
(16684) 1994Q ₁	cClas	6.62 ± 0.12	36.69 ± 6.41	1.13 ± 0.07	0.74 ± 0.10	0.65 ± 0.12	1.29 ± 0.15	2.50 ± 0.20	1.74 ± 0.12	41.892	45.992	3.7	0.0466	43.942	0.36	0.0001281	0.080	3.094	5.912	4, 5, 13, 8
(19255) 1994VK ₈	cClas	7.01 ± 0.16	26.79 ± 7.42	1.01 ± 0.06	0.66 ± 0.10	0.49 ± 0.26	1.15 ± 0.20	—	1.68 ± 0.07	41.673	44.445	1.5	0.0322	43.059	4.22	0.0050888	0.467	2.809	5.158	4, 2, 11, 28
(19299) 1996SZ ₄	Plut	8.18 ± 0.16	13.81 ± 4.73	0.78 ± 0.09	0.52 ± 0.06	0.45 ± 0.15	0.97 ± 0.22	1.54 ± 0.12	1.30 ± 0.10	29.375	50.354	4.7	0.2631	39.865	1.30	0.0012565	0.276	2.968	5.453	7, 3, 5
(19308) 1996TO ₆₆	19:11(b)	4.53 ± 0.04	8.12 ± 8.43	0.68 ± 0.07	0.39 ± 0.15	0.35 ± 0.18	0.74 ± 0.07	1.42 ± 0.10	1.06 ± 0.21	38.382	48.402	27.4	0.1155	43.392	2.22	0.0050824	0.475	2.811	5.212	38, 12, 39, 4, 39, 7, 2, 24, 9
(19521) Chaos	hClas	4.44 ± 0.07	24.58 ± 2.94	0.95 ± 0.05	0.60 ± 0.04	0.64 ± 0.07	1.24 ± 0.06	2.19 ± 0.07	1.56 ± 0.06	41.024	51.092	12.0	0.1093	46.058	1.04	0.0011035	0.235	3.059	5.898	14, 11, 10, 3
(20000) Varuna	hClas	3.34 ± 0.06	24.07 ± 1.91	0.93 ± 0.13	0.60 ± 0.04	0.61 ± 0.02	1.23 ± 0.13	2.15 ± 0.12	1.52 ± 0.04	40.880	45.495	17.1	0.0534	43.188	1.38	0.0020522	0.299	2.983	5.619	11, 27
(20018) 1995QZ ₉	Plut	7.88 ± 0.40	14.08 ± 4.69	0.88 ± 0.04	0.52 ± 0.05	—	1.04 ± 0.13	—	1.40 ± 0.05	33.820	45.426	19.5	0.1465	39.623	1.75	0.0031635	0.365	2.899	5.276	4, 3
(24835) 1995SM ₅₅	hClas(b)	4.35 ± 0.04	1.27 ± 2.91	0.65 ± 0.03	0.37 ± 0.05	0.35 ± 0.05	0.70 ± 0.05	1.35 ± 0.06	1.02 ± 0.05	37.444	46.644	27.0	0.1094	42.044	2.22	0.0050888	0.467	2.809	5.158	4, 2, 11, 28
(24952) 1997Q ₄	Plut	7.38 ± 0.11	6.21 ± 6.38	0.81 ± 0.09	0.42 ± 0.12	0.43 ± 0.08	0.77 ± 0.12	1.53 ± 0.12	1.10 ± 0.10	30.455	48.689	16.6	0.2304	39.572	1.75	0.0028936	0.367	2.900	5.276	4, 7, 14, 5
(24978) 1998HU ₁₅₁	cClas	7.01 ± 0.05	34.99 ± 2.91	1.11 ± 0.03	0.71 ± 0.03	—	1.10 ± 0.11	—	1.82 ± 0.04	41.074	45.271	2.4	0.0486	43.172	0.29	0.0000746	0.064	3.088	5.870	13
(26181) 1996GQ ₂₁	11:2	4.47 ± 0.09	34.74 ± 3.15	1.00 ± 0.06	0.70 ± 0.05	0.69 ± 0.06	1.42 ± 0.07	2.43 ± 0.10	1.69 ± 0.08	38.200	147.441	13.4	0.5885	92.821	1.95	0.0029069	0.632	3.088	6.701	5, 25
(26308) 1998SM ₁₆₅	2:1(b)	5.75 ± 0.12	29.02 ± 4.33	0.98 ± 0.08	0.64 ± 0.07	0.69 ± 0.09	1.34 ± 0.06	2.25 ± 0.09	1.62 ± 0.10	30.107	65.593	13.5	0.3708	47.850	1.89	0.0029334	0.438	2.907	5.586	14, 23, 3
(26375) 1999DE ₃	5:2	4.76 ± 0.07	20.12 ± 2.25	0.93 ± 0.10	0.58 ± 0.04	0.55 ± 0.04	1.16 ± 0.12	2.08 ± 0.11	1.56 ± 0.05	32.229	79.048	7.6	0.4207	55.638	1.76	0.0023039	0.441	2.987	5.974	7, 14, 11, 20, 35, 18
(28978) Ixion	Plut	3.37 ± 0.04	23.01 ± 2.04	0.98 ± 0.18	0.60 ± 0.03	0.58 ± 0.04	1.06 ± 0.24	2.04 ± 0.22	1.63 ± 0.04	29.740	49.006	19.7	0.2447	39.373	2.00	0.0038284	0.416	2.853	5.155	11, 40, 35
(29981) 1999TD ₁₀	SDO	8.68 ± 0.04	13.38 ± 1.78	0.81 ± 0.12	0.51 ± 0.02	0.51 ± 0.05	0.98 ± 0.11	1.82 ± 0.12	1.23 ± 0.03	12.335	185.366	6.0	0.8752	98.851	2.64	0.0049522	0.881	2.049	4.247	14, 11, 35
(31824) Elatus	Cent	10.42 ± 0.11	25.95 ± 3.71	1.03 ± 0.05	0.61 ± 0.05	0.67 ± 0.08	1.28 ± 0.05	2.30 ± 0.08	1.67 ± 0.07	7.290	16.312	5.2	0.3823	11.801	3.40	0.0084471	0.393	3.701	3.212	36, 11
(32532) Thereus	Cent	9.34 ± 0.04	9.48 ± 2.01	0.79 ± 0.07	0.46 ± 0.03	0.48 ± 0.04	0.92 ± 0.08	1.72 ± 0.08	1.19 ± 0.03	8.589	12.753	20.3	0.1951	10.671	3.69	0.0135553	0.399	3.913	3.121	20, 41, 31, 35, 35
(32929) 1995QY ₉	Plut	7.48 ± 0.13	11.04 ± 4.95	0.70 ± 0.12	0.52 ± 0.09	0.40 ± 0.06	0.86 ± 0.09	1.54 ± 0.11	1.16 ± 0.15	29.230	50.467	4.8	0.2665	39.848	1.32	0.0012922	0.280	2.966	5.446	1, 38, 4
(33001) 1997CU ₂₉	cClas	6.17 ± 0.08	28.90 ± 4.47	1.13 ± 0.11	0.66 ± 0.07	0.63 ± 0.09	1.29 ± 0.10	2.43 ± 0.21	1.80 ± 0.12	42.177	45.109	1.5	0.0336	43.643	0.19	0.0000309	0.042	3.096	5.906	7, 17, 6, 8, 10, 3
(33128) 1998BU ₄₈	SDO	6.89 ± 0.13	27.38 ± 6.34	1.04 ± 0.07	0.63 ± 0.10	0.64 ± 0.11	1.25 ± 0.13	2.23 ± 0.10	1.69 ± 0.09	20.522	46.234	14.2	0.3852	33.378	2.36	0.0045888	0.457	2.786	4.687	14, 11, 31
(33340) 1998VG ₄₄	Plut	6.29 ± 0.08	19.02 ± 3.08	0.95 ± 0.05	0.56 ± 0.04	0.60 ± 0.10	1.17 ± 0.11	2.12 ± 0.13	1.51 ± 0.06	29.367	49.786	3.0	0.2580	39.577	1.25	0.0011218	0.263	2.974	5.453	2, 17, 11, 25
(35671) 1998SN ₁₆₅	SDO	5.43 ± 0.07	8.24 ± 4.91	0.75 ± 0.08	0.43 ± 0.06	0.58 ± 0.08	1.06 ± 0.12	1.56 ± 0.12	1.12 ± 0.08	36.278	39.726	4.6	0.0454	38.002						

Table 5 cont'd

Object	Class ^a	H_R^b	Gr^c	$B-V$	$V-R$	$R-I$	$V-I$	$B-I$	$B-R$	q	Q	i	e	a	u_v^d	u^e	e^f	T_N^g	References
(42355) Typhon	SDO(b)	7.36 ± 0.08	13.97 ± 3.01	0.78 ± 0.11	0.52 ± 0.04	0.42 ± 0.12	0.93 ± 0.13	1.71 ± 0.13	1.29 ± 0.07	17.514	58.350	2.4	0.5383	37.932	2.61	0.0048212	0.540	2.684	4.684 19, 20, 35,
(44594) 1999OX ₃	SDO	6.83 ± 0.08	32.26 ± 3.94	1.13 ± 0.07	0.70 ± 0.06	0.62 ± 0.07	1.33 ± 0.08	2.51 ± 0.09	1.84 ± 0.09	17.576	47.072	2.6	0.4562	32.324	2.40	0.0040896	0.459	2.774	4.592 3, 31, 17, 14, 11, 22, 31, 23, 24, 27
(47171) 1999TC ₃₆	Plut(b)	4.85 ± 0.05	32.06 ± 2.34	0.99 ± 0.13	0.69 ± 0.03	0.62 ± 0.05	1.30 ± 0.13	2.35 ± 0.14	1.74 ± 0.05	30.556	48.674	8.4	0.2287	39.615	1.28	0.0013649	0.271	2.970	5.446 2, 17, 14, 20, 44, 35
(47932) 2000GN ₁₇₁	Plut	5.67 ± 0.09	23.88 ± 2.88	0.96 ± 0.06	0.60 ± 0.04	0.62 ± 0.05	1.14 ± 0.17	2.16 ± 0.09	1.56 ± 0.07	28.280	50.231	10.8	0.2796	39.255	1.60	0.0021425	0.337	2.921	5.313 5, 25
(48639) 1995TL ₈	dTNO(b)	4.66 ± 0.09	23.17 ± 6.08	1.01 ± 0.16	0.62 ± 0.09	0.55 ± 0.12	1.14 ± 0.21	2.25 ± 0.32	1.69 ± 0.22	40.202	65.463	0.2	0.2391	52.832	0.98	0.0006765	0.239	3.143	6.287 14, 11, 24
(49036) Pelion	Cent	10.15 ± 0.11	18.99 ± 6.95	0.74 ± 0.08	0.56 ± 0.09	0.58 ± 0.21	0.84 ± 0.14	1.61 ± 0.17	1.25 ± 0.10	17.325	22.710	9.3	0.1345	20.018	1.41	0.0018835	0.211	3.098	4.095 31, 3, 31, 5, 11
(50000) Quaoar	hClas(b)	2.22 ± 0.03	63.39 ± 1.34	0.94 ± 0.11	0.65 ± 0.02	1.28 ± 0.02	1.33 ± 0.14	2.25 ± 0.13	1.59 ± 0.02	41.607	44.752	8.0	0.0364	43.179	0.65	0.0004670	0.144	3.068	5.822 35, 26, 35, 20
(52747) 1998HM ₁₅₁	cClas	7.42 ± 0.10	24.89 ± 4.69	0.93 ± 0.09	0.62 ± 0.05	—	—	—	1.55 ± 0.10	41.704	46.760	0.5	0.0572	44.232	0.26	0.0000482	0.058	3.101	5.939 13
(52872) Okyho	Cent	10.78 ± 0.08	11.27 ± 3.22	0.74 ± 0.07	0.48 ± 0.05	0.48 ± 0.06	0.96 ± 0.06	1.72 ± 0.08	1.24 ± 0.09	5.781	10.891	15.7	0.3065	8.336	4.23	0.0157838	0.408	4.572	2.944 31, 17, 14, 44, 45
(52975) Cyllarus	Cent	8.62 ± 0.10	32.34 ± 5.23	1.10 ± 0.10	0.68 ± 0.09	0.67 ± 0.09	1.34 ± 0.10	2.50 ± 0.13	1.80 ± 0.10	16.338	36.273	12.6	0.3789	26.305	2.54	0.0052272	0.437	2.832	4.259 2, 31, 11, 31, 20
(53311) Deucalion	cClas	6.67 ± 0.06	30.38 ± 20.84	1.36 ± 0.16	0.67 ± 0.23	—	—	—	2.03 ± 0.16	41.339	46.596	0.4	0.0598	43.968	0.27	0.0000517	0.060	3.098	5.922 6
(54598) Bienor	Cent	7.72 ± 0.08	9.06 ± 3.36	0.74 ± 0.14	0.48 ± 0.05	0.40 ± 0.08	0.85 ± 0.14	1.60 ± 0.15	1.16 ± 0.07	13.213	19.886	20.7	0.2016	16.550	3.03	0.0091145	0.408	3.176	3.582 14, 11, 20, 31, 23
(55565) 2002AW ₁₉₇	hClas	3.16 ± 0.06	23.45 ± 3.25	0.88 ± 0.15	0.59 ± 0.04	0.95 ± 0.30	1.21 ± 0.17	2.05 ± 0.17	1.50 ± 0.04	41.312	53.382	24.3	0.1275	47.347	1.92	0.0038014	0.431	2.903	5.562 35, 46, 26, 25
(55576) Amycus	Cent	7.80 ± 0.04	35.81 ± 2.84	1.11 ± 0.03	0.70 ± 0.03	0.76 ± 0.10	1.47 ± 0.13	2.73 ± 0.24	1.81 ± 0.04	15.166	34.750	13.4	0.3923	24.958	2.72	0.0059909	0.455	2.836	4.129 19, 31, 26, 46, 25, 27
(55636) 2002TX ₃₀₀	hClas(b)	3.29 ± 0.05	-0.07 ± 1.39	0.70 ± 0.09	0.36 ± 0.02	0.33 ± 0.03	0.68 ± 0.12	1.39 ± 0.12	1.01 ± 0.03	38.102	48.720	25.9	0.1223	43.411	2.11	0.0046004	0.453	2.839	5.279 20, 22, 35, 28
(55637) 2002UX ₂₅	SDO(b)	3.48 ± 0.08	19.31 ± 2.44	0.95 ± 0.13	0.56 ± 0.04	0.59 ± 0.05	1.05 ± 0.15	2.00 ± 0.14	1.50 ± 0.05	36.689	49.110	19.4	0.1448	42.899	1.67	0.0028799	0.363	2.930	5.480 35, 15, 25
(55638) 2002VE ₉₅	Plut	5.14 ± 0.06	37.77 ± 3.71	1.07 ± 0.14	0.72 ± 0.05	0.76 ± 0.12	1.38 ± 0.15	2.47 ± 0.13	1.79 ± 0.04	28.001	51.242	16.3	0.2933	39.621	1.93	0.0033497	0.406	2.865	5.195 25, 35, 27
(58534) Logos	cClas(b)	6.76 ± 0.18	34.12 ± 7.19	0.99 ± 0.13	0.73 ± 0.12	0.60 ± 0.12	1.20 ± 0.13	2.25 ± 0.18	1.65 ± 0.15	39.735	50.867	2.9	0.1229	45.301	0.59	0.0002647	0.133	3.097	5.964 7, 2, 4, 10, 8
(59358) 1999CL ₁₅₈	SDO	6.65 ± 0.09	4.54 ± 2.77	0.80 ± 0.06	0.39 ± 0.04	0.47 ± 0.06	0.86 ± 0.06	1.66 ± 0.09	1.19 ± 0.07	32.839	50.459	10.0	0.2115	41.649	1.26	0.0013977	0.274	2.987	5.571 11
(60454) 2000CH ₁₀₅	cClas	6.37 ± 0.08	26.14 ± 5.75	1.02 ± 0.09	0.64 ± 0.10	0.58 ± 0.10	1.23 ± 0.10	2.23 ± 0.10	1.70 ± 0.08	40.647	48.067	1.2	0.0836	44.357	0.38	0.0001078	0.086	3.098	5.935 19, 8
(60458) 2000CM ₁₁₄	SDO(b)	6.95 ± 0.04	14.16 ± 1.89	0.73 ± 0.02	0.50 ± 0.02	0.66 ± 0.09	0.87 ± 0.10	—	1.24 ± 0.04	35.536	83.826	19.7	0.4046	59.681	2.04	0.0036132	0.526	2.930	5.920 20, 15
(60558) Echeclus	Cent	9.68 ± 0.09	12.93 ± 4.19	0.85 ± 0.05	0.51 ± 0.06	0.49 ± 0.10	0.95 ± 0.10	1.76 ± 0.12	1.38 ± 0.07	5.816	15.579	4.3	0.4563	10.697	4.21	0.0127027	0.463	3.869	3.031 5, 31, 25
(60608) 2000EE ₁₇₃	SDO	8.03 ± 0.11	12.78 ± 2.20	0.67 ± 0.03	0.49 ± 0.03	0.54 ± 0.06	1.02 ± 0.06	1.74 ± 0.11	1.16 ± 0.03	22.576	76.086	5.9	0.5424	49.331	2.34	0.0039446	0.552	2.750	5.251 5, 8, 20
(60620) 2000FD ₈	7:4	6.35 ± 0.06	30.23 ± 4.54	1.15 ± 0.11	0.66 ± 0.08	0.65 ± 0.07	1.32 ± 0.10	2.44 ± 0.14	1.81 ± 0.11	34.124	57.209	19.5	0.2185	43.666	1.82	0.0032402	0.399	2.905	5.448 5, 8, 19
(60621) 2000FE ₈	5:2(b)	6.51 ± 0.06	11.21 ± 1.71	0.75 ± 0.02	0.48 ± 0.02	0.50 ± 0.03	0.98 ± 0.05	1.73 ± 0.09	1.23 ± 0.03	32.994	73.643	5.9	0.4036	55.318	1.67	0.0020292	0.416	3.013	6.029 11, 20
(63252) 2001BL ₄₁	Cent	11.29 ± 0.07	11.69 ± 3.77	0.72 ± 0.05	0.50 ± 0.04	0.38 ± 0.16	1.01 ± 0.08	1.75 ± 0.08	1.20 ± 0.04	6.834	12.607	12.5	0.2970	9.721	3.51	0.0104763	0.367	4.154	3.084 31, 19, 20
(65489) Ceto	SDO(b)	6.21 ± 0.06	18.61 ± 2.91	0.86 ± 0.03	0.56 ± 0.03	—	—	—	1.42 ± 0.04	17.760	184.198	22.3	0.8241	100.979	2.70	0.0056291	0.907	2.219	4.669 20
(66452) 1999OF ₄	cClas	6.25 ± 0.09	35.23 ± 4.18	1.03 ± 0.11	0.74 ± 0.07	0.60 ± 0.08	1.26 ± 0.14	2.43 ± 0.12	1.83 ± 0.10	42.071	47.731	2.7	0.0630	44.901	0.35	0.0001033	0.078	3.106	5.973 19
(66652) Borasiri	cClas(b)	5.42 ± 0.05	34.20 ± 3.90	0.86 ± 0.09	0.71 ± 0.06	0.64 ± 0.07	1.29 ± 0.07	—	1.61 ± 0.05	39.994	47.575	0.6	0.0866	43.785	0.39	0.0001092	0.087	3.091	5.898 17, 14, 21
(69866) 1998WW ₂₄	Plut	7.96 ± 0.10	14.56 ± 5.09	0.76 ± 0.11	0.47 ± 0.08	0.66 ± 0.11	1.13 ± 0.09	1.88 ± 0.13	1.24 ± 0.15	30.644	48.774	13.9	0.2283	39.709	1.57	0.0022755	0.332	2.929	5.352 11, 19
(69877) 1998WA ₂₅	cClas	—	—	—	—	—	1.08 ± 0.11	—	—	41.771	43.746	1.0	0.0231	42.759	0.13	0.0000156	0.029	3.087	5.852 8
(69888) 1998WA ₃₁	5:2	7.30 ± 0.15	16.79 ± 7.68	0.79 ± 0.13	0.54 ± 0.11	0.53 ± 0.18	1.08 ± 0.18	1.83 ± 0.19	1.41 ± 0.13	31.682	79.362	9.5	0.4294	55.522	1.84	0.0025611	0.460	2.963	5.914 8, 19
(69990) 1998WU ₃₁	Plut	7.98 ± 0.20	26.01 ± 4.86	0.72 ± 0.09	0.72 ± 0.09	0.41 ± 0.07	1.13 ± 0.11	1.64 ± 0.11	1.23 ± 0.09	32.006	46.999	6.6	0.1898	39.503	1.05	0.0009006	0.222	2.977	5.507 19
(73480) 2002PN ₃₄	SDO	8.48 ± 0.05	12.37 ± 4.87	0.83 ± 0.12	0.50 ± 0.05	—	0.98 ± 0.16	1.82 ± 0.14	1.28 ± 0.02	13.365	48.624	16.6	0.5688	30.995	3.41	0.0091683	0.637	2.970	4.015 20, 35, 45
(79360) Sila-Nunam	cClas(b)	5.07 ± 0.09	28.39 ± 3.34	1.08 ± 0.07	0.67 ± 0.05	0.59 ± 0.06	1.27 ± 0.07	2.37 ± 0.11	1.75 ± 0.08	43.447	44.675	2.2	0.0139	44.061	0.19	0.0000374	0.041	3.101	5.933 7, 2, 9, 10
(79969) 1999CP ₁₃₃	5:4	7.24 ± 0.04	8.27 ± 2.46	0.78 ± 0.06	0.45 ± 0.04	0.45 ± 0.03	0.90 ± 0.05	1.68 ± 0.05	1.23 ± 0.04	31.948	37.930	3.2	0.0854	34.930	0.51	0.0002181	0.102	3.005	5.304 18
(79978) 1999CC ₁₅₈	12:5	5.42 ± 0.08	23.34 ± 3.50	1.01 ± 0.07	0.60 ± 0.06	0.60 ± 0.06	1.22 ± 0.07	2.26 ± 0.10	1.64 ± 0.09	39.063	69.390	18.7	0.2796	54.226	1.74	0.0028021	0.426	2.997	5.967 14, 11, 25, 18
(79983) 1999DF ₉	hClas	5.80 ± 0.11	34.29 ± 3.22	0.92 ± 0.06	0.71 ± 0.05	0.65 ± 0.06	1.36 ± 0.06	2.28 ± 0.10	1.63 ± 0.08	39.752	53.254	9.8	0.1452	46.503	0.98	0.0009072	0.224	3.071	5.941 11
(80806) 2000CM ₁₀₅	cClas(b)	6.30 ± 0.03	51.15 ± 29.94	1.14 ± 0.23	0.84 ± 0.33	—	1.34 ± 0.10	—	1.98 ± 0.23	39.427	45.196	3.8	0.0682	42.311	0.43	0.0001702	0.095	3.073	5.801 6
(82075) 2000YW ₁₃₄	8:3(b)	4.43 ± 0.06	14.45 ± 3.51	0.92 ± 0.05	0.48 ± 0.06	0.60 ± 0.06	1.08 ± 0.07	1.99 ± 0.08	1.42 ± 0.08	41.130	75.402	19.8	0.2941	58.266	1.77	0.0028931	0.448	3.020	6.109 19, 22, 15, 24, 20
(82155) 2001FZ ₁₇₃	SDO	5.82 ± 0.03	17.27 ± 1.84	0.86 ± 0.03	0.56 ± 0.02	0.50 ± 0.05	1.08 ± 0.06	1.96 ± 0.08	1.42 ± 0.03	32.396	137.849	12.7	0.6194	85.123	2.12	0.0033866	0.657	2.930	6.256 19, 20
(82158) 2001FP ₁₈₅	SDO	5.94 ± 0.05	18.28 ± 2.69	0.81 ± 0.05	0.58 ± 0.03	0.42 ± 0.08	1.01 ± 0.08	1.89 ± 0.14	1.40 ± 0.06	34.231	398.411	30.8	0.8418	216.321	2.02	0.0032595	0.985	2.626	6.004 19, 22, 20
(83982) Crantor	Cent	8.71 ± 0.06	43.27 ± 3.40	1.10 ± 0.04	0.76 ± 0.04	0.92 ± 0.19	1.65 ± 0.18	2.85 ± 0.28	1.86 ± 0.04	14.025	24.676	12.8	0.2752	19.351	2.40	0.0049784	0.353	3.058	3.885 19, 31, 20, 46, 26
(84522) 2002TC ₃₀₂	5:2	3.68 ± 0.07	29.33 ± 2.40	1.09 ± 0.06	0.65 ± 0.04	0.66 ± 0.04	1.30 ± 0.06	2.42 ± 0.06	1.74 ± 0.05	39.108	72.283	35.0	0.2978	55.695	2.71	0.0069045	0.646	2.668	5.210 15, 25, 24
(84719) 2002VR ₁₂₈	Plut	5.00 ± 0.04	22.76 ± 2.06	0.94 ± 0.03	0.60 ± 0.02	—	—	—	1.54 ± 0.04	29.176	50.011	14.0	0.2631	39.594	1.70	0.0025714	0.357	2.908	5.296 25
(84922) 2003V ₅₂	Plut	3.7																	

Table 5 cont'd

Object	Class ^a	H_R^b	Gr^c	$B-V$	$V-R$	$R-I$	$V-I$	$B-I$	$B-R$	q	Q	i	e	u_v^d	u^e	e^f	T_N^g	T_h^h	References
(85633) 1998KR ₆₅	cClass	6.60 ± 0.07	32.04 ± 4.24	1.09 ± 0.08	0.63 ± 0.07	0.79 ± 0.08	1.33 ± 0.09	2.30 ± 0.10	1.73 ± 0.14	41.972	44.911	1.2	0.0338	43.441	0.18	0.0000264	0.040	3.094	5.894 13, 8, 5
(86047) 1999OY ₃	SDO(h)	-2.62 ± 3.39	0.72 ± 0.03	0.33 ± 0.05	0.28 ± 0.08	0.57 ± 0.07	1.30 ± 0.09	1.05 ± 0.05	0.60 ± 0.05	36.401	51.117	24.3	0.1682	43.759	2.05	0.00042644	0.444	2.855	5.331 3, 5, 11
(86177) 1999RX ₂₁₅	SDO	10.27 ± 6.57	0.72 ± 0.12	0.44 ± 0.09	0.63 ± 0.18	0.70 ± 0.25	1.78 ± 0.34	1.15 ± 0.18	34.501	56.169	22.2	0.2390	45.335	2.01	0.0039357	0.447	2.871	5.422 17, 5	
(87269) 2000OQ ₆₇	SDO	27.14 ± 4.40	1.08 ± 0.09	0.65 ± 0.08	0.59 ± 0.06	1.24 ± 0.07	2.30 ± 0.04	1.70 ± 0.09	20.794	119.714	20.1	0.9659	60.254	1.24	0.0011503	1.025	2.240	5.274 20, 8, 24	
(87555) 2000QB ₂₄₃	SDO	8.44 ± 0.12	6.81 ± 6.12	0.76 ± 0.09	0.40 ± 0.08	0.69 ± 0.22	0.94 ± 0.24	1.59 ± 0.09	15.285	54.442	6.8	0.5616	34.864	2.89	0.0006544	0.574	2.632	4.403 19, 23	
(88268) 2001KK ₇₆	cClass	—	47.24 ± 4.69	1.03 ± 0.06	0.81 ± 0.05	—	—	1.84 ± 0.08	41.697	42.958	1.9	0.0149	42.327	0.17	0.0000289	0.036	3.082	5.823 21	
(88269) 2001KF ₇₇	Cent	10.04 ± 0.02	37.35 ± 1.30	1.08 ± 0.04	0.73 ± 0.01	—	—	1.81 ± 0.04	19.779	32.001	4.4	0.2360	25.890	1.45	0.0015681	0.248	2.960	4.524 20	
(88611) Teharonhiawako	cClass(b)	—	—	—	—	—	0.82 ± 0.04	—	42.802	45.195	2.6	0.0272	43.998	0.24	0.0000568	0.053	3.999	5.926 49	
(90377) Sedna	dTNO	1.11 ± 0.09	37.31 ± 7.28	1.17 ± 0.12	0.73 ± 0.12	0.68 ± 0.13	1.39 ± 0.15	2.51 ± 0.14	1.87 ± 0.12	76.223	1010.285	11.9	0.8597	543.254	1.13	0.0009289	0.884	4.304	10.223 35, 25, 27, 24
(90482) Oresus	Plut(b)	1.99 ± 0.05	1.92 ± 2.47	0.62 ± 0.08	0.38 ± 0.04	0.37 ± 0.05	0.75 ± 0.10	1.34 ± 0.09	1.04 ± 0.04	30.649	48.087	20.5	0.2215	39.368	2.00	0.0039069	0.415	2.853	5.156 35, 47, 35, 25, 45
(90568) 2004GV ₉	hClass	3.79 ± 0.08	14.61 ± 2.91	0.90 ± 0.06	0.52 ± 0.03	—	1.15 ± 0.03	1.99 ± 0.04	1.47 ± 0.04	38.679	45.085	22.0	0.0765	41.882	1.80	0.0034450	0.383	2.900	5.369 25, 28
(91133) 1998HK ₁₅₁	Plut	6.94 ± 0.08	11.23 ± 3.75	0.72 ± 0.05	0.52 ± 0.06	0.39 ± 0.06	0.91 ± 0.07	1.60 ± 0.08	1.24 ± 0.06	30.359	48.056	5.9	0.2257	39.208	1.18	0.0010857	0.248	2.980	5.452 2, 17, 11, 8
(91205) 1998US ₁₃	Plut	7.85 ± 0.05	7.54 ± 5.23	0.69 ± 0.10	0.49 ± 0.09	0.32 ± 0.07	0.82 ± 0.12	1.51 ± 0.12	1.19 ± 0.10	34.194	44.813	10.6	0.1344	39.503	1.08	0.0011405	0.228	2.994	5.499 19
(91554) 1999RZ ₂₁₅	SDO	19.40 ± 5.44	0.77 ± 0.10	0.57 ± 0.09	0.54 ± 0.09	1.11 ± 0.13	1.89 ± 0.16	1.35 ± 0.13	30.989	172.672	25.5	0.6957	101.831	2.44	0.0047930	0.818	2.681	5.787 5	
(95625) 2002GX ₃₂	7:3	7.47 ± 0.06	37.37 ± 6.12	—	0.77 ± 0.09	0.56 ± 0.13	1.33 ± 0.19	—	—	33.079	72.609	13.9	0.3740	52.844	1.83	0.0027543	0.445	2.955	5.835 1
(95626) 2002GZ ₅₂	Cent	6.61 ± 0.13	21.51 ± 3.72	0.71 ± 0.13	0.58 ± 0.05	0.61 ± 0.09	1.07 ± 0.13	1.74 ± 0.17	1.20 ± 0.07	18.007	28.007	15.0	0.2173	23.007	2.11	0.0042076	0.338	2.956	4.191 22, 31, 20, 35, 26, 35
(118228) 1996TQ ₆₆	Plut	7.24 ± 0.20	34.22 ± 6.08	1.19 ± 0.12	0.67 ± 0.10	0.75 ± 0.11	1.47 ± 0.15	2.66 ± 0.19	1.88 ± 0.14	34.638	44.664	14.7	0.1264	39.651	1.35	0.0018674	0.283	2.962	5.430 4, 7, 8, 9
(118378) 1999HT ₁₁	7:4	6.85 ± 0.07	31.69 ± 4.07	1.15 ± 0.10	0.67 ± 0.09	0.66 ± 0.03	1.33 ± 0.05	2.48 ± 0.06	1.82 ± 0.07	38.789	48.457	5.1	0.1108	43.623	0.64	0.0003542	0.142	3.074	5.852 6, 18
(118379) 1999HC ₁₂	SDO	—	8.01 ± 8.06	0.89 ± 0.16	0.49 ± 0.14	0.34 ± 0.12	1.02 ± 0.22	1.73 ± 0.25	1.38 ± 0.21	34.575	55.799	15.4	0.2348	45.187	1.58	0.0023181	0.354	2.963	5.639 5
(118702) 2000OM ₆₇	SdTNO	13.46 ± 2.46	0.82 ± 0.06	0.47 ± 0.04	0.59 ± 0.04	1.06 ± 0.06	1.88 ± 0.06	1.29 ± 0.04	30.256	157.682	23.4	0.6013	98.469	2.18	0.0038941	0.720	2.960	6.434 24	
(119068) 2001KC ₇₇	5:2	6.83 ± 0.03	18.61 ± 1.30	0.91 ± 0.01	0.56 ± 0.01	—	—	1.47 ± 0.01	35.399	74.238	12.9	0.3542	54.819	1.69	0.0023431	0.419	3.010	6.012 20	
(119070) 2001KP ₇₇	7:4	6.88 ± 0.30	20.48 ± 7.24	1.10 ± 0.29	0.59 ± 0.10	0.52 ± 0.19	1.07 ± 0.17	2.18 ± 0.43	1.72 ± 0.32	36.004	50.980	3.3	0.1722	43.492	0.82	0.0005029	0.182	3.057	5.806 19, 22
(119315) 2001SD ₇₃	Cent	8.84 ± 0.07	10.81 ± 2.65	0.67 ± 0.02	0.46 ± 0.03	0.57 ± 0.09	1.07 ± 0.13	—	1.13 ± 0.02	14.473	20.567	17.4	0.1739	17.520	2.49	0.0001987	0.346	3.151	3.745 20, 15
(119473) 2001UO ₁₈	Plut	7.80 ± 0.51	46.23 ± 19.58	1.27 ± 0.30	0.80 ± 0.21	—	—	2.08 ± 0.38	28.274	51.587	3.7	0.2919	39.930	1.41	0.0014361	0.299	2.953	5.419 22	
(119878) 2002CY ₂₂₄	12:5	5.89 ± 0.04	30.91 ± 2.70	1.12 ± 0.10	0.67 ± 0.04	0.65 ± 0.04	1.32 ± 0.06	2.45 ± 0.12	1.81 ± 0.10	35.236	73.190	15.7	0.3500	54.213	1.82	0.0027651	0.443	2.976	5.917 15, 18
(119951) 2002KX ₁₄	SDO	4.35 ± 0.12	23.82 ± 2.06	1.05 ± 0.03	0.61 ± 0.02	—	1.21 ± 0.16	—	1.66 ± 0.04	36.918	40.400	0.4	0.0451	38.662	0.20	0.000342	0.046	3.043	5.581 25, 29
(119956) 2002PA ₁₄₉	7:4	6.21 ± 0.03	25.38 ± 1.92	1.14 ± 0.04	0.65 ± 0.04	0.56 ± 0.02	1.21 ± 0.04	2.35 ± 0.03	1.79 ± 0.02	36.129	51.143	4.1	0.1720	43.636	0.84	0.0005383	0.186	3.057	5.810 18
(119979) 2002WC ₁₉	2:1(b)	4.25 ± 0.01	29.36 ± 0.94	1.14 ± 0.02	0.65 ± 0.01	1.30 ± 0.04	2.44 ± 0.02	1.79 ± 0.02	35.466	60.971	9.2	0.2645	48.218	1.32	0.0014333	0.309	3.035	5.904 18	
(120061) 2003CO ₁	Cent	9.14 ± 0.14	9.93 ± 4.38	0.74 ± 0.03	0.48 ± 0.05	0.38 ± 0.17	0.81 ± 0.17	—	1.24 ± 0.04	10.906	30.453	19.8	0.4726	20.680	3.83	0.0122807	0.581	2.830	3.558 20, 27
(120132) 2003FY ₁₂₈	SDO	4.49 ± 0.05	21.48 ± 3.46	1.05 ± 0.03	0.60 ± 0.05	0.54 ± 0.08	1.14 ± 0.09	2.20 ± 0.04	1.65 ± 0.02	36.980	61.840	11.8	0.2516	49.410	1.37	0.0016427	0.324	3.038	5.945 27, 24
(120178) 2003OR ₃₂	hClass(h)	3.74 ± 0.05	2.40 ± 3.75	0.70 ± 0.05	0.39 ± 0.06	0.36 ± 0.07	0.76 ± 0.06	1.47 ± 0.07	—	38.650	47.601	27.2	0.1038	43.126	2.20	0.0049859	0.468	2.817	5.216 27, 28
(120181) 2003UR ₉₂	SDO	7.09 ± 0.10	28.17 ± 4.69	1.04 ± 0.06	0.65 ± 0.05	—	—	1.69 ± 0.08	26.768	38.252	2.7	0.1766	32.510	2.85	0.0006689	0.183	2.970	5.075 25	
(120216) 2004EW ₉₅	Plut	6.31 ± 0.05	1.71 ± 2.06	0.70 ± 0.02	0.38 ± 0.02	—	—	1.08 ± 0.03	26.976	51.657	29.3	0.3139	39.316	2.85	0.0076582	0.581	2.658	4.684 25	
(120347) Salacia	hClass(b)	3.82 ± 0.01	—	—	—	—	0.87 ± 0.01	—	37.687	46.464	37.687	23.9	0.1043	42.075	1.98	0.0040765	0.419	2.865	5.293 49, 16
(120348) 2004TY ₃₆₄	SDO	4.11 ± 0.04	20.78 ± 3.17	0.92 ± 0.18	0.60 ± 0.05	0.52 ± 0.06	1.03 ± 0.18	1.95 ± 0.17	—	36.437	41.611	24.8	0.0663	39.024	2.99	0.0045940	0.425	2.834	5.093 35, 27
(121725) 1999XX ₁₄₃	Cent	8.59 ± 0.10	30.41 ± 7.61	1.11 ± 0.15	0.65 ± 0.12	0.69 ± 0.14	1.31 ± 0.16	2.47 ± 0.48	1.73 ± 0.14	9.687	26.257	6.8	0.4610	17.972	3.34	0.0081625	0.476	3.036	3.565 31, 11, 19
(123509) 2000WK ₁₈₃	cClass(b)	—	—	—	—	—	1.30 ± 0.05	—	42.437	46.910	2.0	0.0501	44.673	0.27	0.0000615	0.061	3.106	5.966 49	
(126154) 2001YH ₄₀	5:3	5.38 ± 0.07	17.01 ± 3.18	0.97 ± 0.08	0.56 ± 0.07	0.51 ± 0.02	1.07 ± 0.07	2.04 ± 0.04	1.53 ± 0.03	36.383	48.798	11.1	0.1457	42.590	1.10	0.0011760	0.241	3.017	5.678 18
(126619) 2002CX ₁₅₄	11:3	7.24 ± 0.09	20.39 ± 4.61	0.95 ± 0.11	0.57 ± 0.07	0.57 ± 0.08	1.14 ± 0.11	2.12 ± 0.15	1.54 ± 0.10	37.960	105.208	16.0	0.4697	71.584	1.92	0.0029838	0.544	3.039	6.369 15, 18
(127546) 2002XU ₉₃	cClass	6.05 ± 0.07	20.07 ± 4.44	—	0.51 ± 0.08	0.69 ± 0.06	1.20 ± 0.10	—	38.011	56.127	77.9	0.6875	67.167	5.53	0.0186306	1.195	0.903	1.173 24	
(127871) 2003FC ₁₂₈	5:4	6.91 ± 0.03	36.85 ± 1.92	1.20 ± 0.05	0.69 ± 0.04	0.74 ± 0.02	1.43 ± 0.04	2.63 ± 0.04	1.89 ± 0.03	31.917	38.062	2.4	0.0878	34.990	0.49	0.0001867	0.097	3.007	5.311 18
(129772) 1999HR ₁₁	7:4	7.18 ± 0.15	26.14 ± 5.45	0.92 ± 0.12	0.53 ± 0.10	0.80 ± 0.07	1.33 ± 0.12	2.25 ± 0.17	1.45 ± 0.16	42.165	45.032	3.3	0.0329	43.599	0.30	0.0000918	0.066	3.093	5.896 17
(130391) 2000JG ₈₁	2:1	7.74 ± 0.06	8.20 ± 5.50	0.79 ± 0.08	0.42 ± 0.09	0.51 ± 0.08	0.91 ± 0.13	1.77 ± 0.06	1.32 ± 0.10	34.125	60.398	23.5	0.2779	47.262	2.14	0.0043818	0.486	2.845	5.421 4, 18
(131696) 2001XT ₂₅₄	7:3	7.29 ± 0.06	9.40 ± 3.40	0.85 ± 0.08	0.49 ± 0.06	0.40 ± 0.05	0.89 ± 0.08	1.74 ± 0.07	1.34 ± 0.05	35.898	70.816	0.5	0.3272	53.357	1.33	0.0012557	0.327	3.081	6.149 18
(131697) 2001XH ₂₅₅	5:4	8.18 ± 0.07	14.39 ± 4.14	0.85 ± 0.09	0.49 ± 0.07	0.57 ± 0.06	1.06 ± 0.09	1.91 ± 0.09	1.34 ± 0.06	32.413	37.783	2.9	0.0765	35.098	0.46	0.0001747	0.091	3.009	5.321 18
(134340) Pluto	Plut(b)	-1.33 ± 0.01	8.31 ± 0.82	0.86 ± 0.01	0.48 ± 0.01	0.40 ± 0.01	1.74 ± 0.02	1.34 ± 0.01	1.72 ± 0.08	42.011	43.623	17.2	0.2470	39.348	1.84	0.0031834	0.385	2.882	5.224 7
(134860) 2000OI ₆₇	cClass(b)	6.00 ± 0.12	29.15 ± 3.37	1.05 ± 0.06	0.67 ± 0.05	0.60 ± 0.07	1.20 ± 0.10	2.32 ± 0.10	1.72 ± 0.08	42.011	43.623	11.1	0.0188	42.817	0.12	0.0000140	0.027	3.088	5.857

Table 5 cont'd

Object	Class ^a	H_R^b	Gr^c	$B-V$	$V-R$	$R-I$	$V-I$	$B-I$	$B-R$	q	Q	i	e	a	u_v^d	u^e	T_N^g	T^h	References
(143707) 2003UY ₁₁₇	5:2	5.55 ± 0.01	20.28 ± 0.94	0.97 ± 0.02	0.56 ± 0.01	0.59 ± 0.01	1.15 ± 0.02	1.12 ± 0.01	1.53 ± 0.01	32.489	79.484	7.5	0.4197	55.986	1.75	0.0022729	0.440	2.993	5.996 18
(144897) 2004UX ₁₀	hClas	4.21 ± 0.09	20.23 ± 4.40	0.95 ± 0.02	0.58 ± 0.05	—	—	—	1.53 ± 0.02	37.598	40.720	9.5	0.0399	39.159	0.81	0.0007260	0.170	3.017	5.539 29, 27
(145451) 2005RM ₄₃	SDO	4.25 ± 0.05	3.01 ± 4.89	0.59 ± 0.04	0.40 ± 0.05	—	0.73 ± 0.04	1.32 ± 0.05	—	35.123	148.950	28.7	0.6184	92.036	2.47	0.0051049	0.783	2.738	5.854 27, 28
(145453) 2005RR ₄₃	hClas(h)	3.85 ± 0.04	3.89 ± 4.89	0.79 ± 0.08	0.41 ± 0.05	—	0.69 ± 0.08	1.48 ± 0.11	—	37.318	49.609	28.5	0.1414	43.463	2.33	0.0055111	0.497	2.785	5.151 27, 28
(145480) 2005TB ₁₉₀	dTNO	—	18.63 ± 1.89	0.98 ± 0.04	0.56 ± 0.03	0.55 ± 0.03	1.11 ± 0.04	2.09 ± 0.04	1.54 ± 0.03	46.191	105.344	26.5	0.3904	75.767	2.07	0.0038799	0.593	3.013	6.358 24
(148209) 2000CR ₁₀₅	dTNO	6.19 ± 0.07	14.91 ± 5.95	0.77 ± 0.08	0.47 ± 0.08	0.73 ± 0.17	1.10 ± 0.13	1.85 ± 0.13	1.27 ± 0.07	44.126	413.100	22.7	0.8070	228.613	1.77	0.0024329	0.895	3.136	7.244 20, 15, 24
(148780) Altjira	hClas(b)	5.88 ± 0.32	33.03 ± 8.07	0.91 ± 0.13	0.73 ± 0.11	0.48 ± 0.23	1.23 ± 0.16	2.29 ± 0.21	1.64 ± 0.17	41.886	47.080	5.2	0.0584	44.483	0.48	0.0002326	0.108	3.094	5.931 22
(149349) 2002VA ₁₃₁	5:3	6.42 ± 0.05	40.19 ± 2.66	1.33 ± 0.06	0.77 ± 0.05	0.67 ± 0.03	1.44 ± 0.06	2.77 ± 0.04	2.10 ± 0.03	32.186	52.895	7.1	0.2434	42.540	1.24	0.0012251	0.273	2.997	5.627 18
(149560) 2003QZ ₆₁	SDO	8.30 ± 0.03	—	—	—	—	—	—	1.30 ± 0.05	21.761	61.186	34.8	0.4753	41.473	3.57	0.0112771	0.743	2.421	4.202 50
(160147) 2001KN ₇₆	7:4	6.62 ± 0.04	24.55 ± 2.46	1.05 ± 0.06	0.60 ± 0.04	0.63 ± 0.03	1.23 ± 0.05	2.28 ± 0.05	1.65 ± 0.04	39.855	47.289	2.6	0.0853	43.572	0.44	0.0001533	0.097	3.086	5.880 18
(168700) 2000GE ₄₇	Plut	8.01 ± 0.11	16.97 ± 7.13	—	0.55 ± 0.12	0.52 ± 0.11	1.07 ± 0.11	—	—	30.264	48.312	5.0	0.2297	39.288	1.17	0.0010319	0.246	2.982	5.461 8
(168703) 2000GP ₈₃	SDO	5.80 ± 0.06	8.18 ± 3.45	0.77 ± 0.04	0.45 ± 0.05	0.46 ± 0.07	0.91 ± 0.08	1.59 ± 0.07	1.16 ± 0.06	36.806	42.587	4.9	0.0728	39.696	0.53	0.0002686	0.112	3.041	5.620 11, 8, 19
(174567) 2003MW ₁₂	hClas(b)	3.21 ± 0.10	17.11 ± 7.64	—	0.58 ± 0.11	0.42 ± 0.17	1.00 ± 0.17	—	—	39.052	52.114	21.5	0.1433	45.583	0.77	0.0032322	0.394	2.927	5.564 27
(181708) 1993FW	hClas	6.58 ± 0.10	22.47 ± 4.11	0.97 ± 0.08	0.63 ± 0.06	0.45 ± 0.11	1.06 ± 0.14	2.06 ± 0.12	1.62 ± 0.11	41.636	45.656	7.8	0.0461	43.646	0.64	0.0004476	0.143	3.074	5.853 1, 13, 8, 10
(181855) 1998WT ₃₁	hClas	7.44 ± 0.08	9.74 ± 7.77	0.75 ± 0.14	0.48 ± 0.12	0.42 ± 0.16	0.92 ± 0.22	1.68 ± 0.21	1.25 ± 0.14	37.593	55.055	28.6	0.1885	46.324	2.34	0.0054411	0.515	2.789	5.255 19, 16
(181867) 1999CV ₁₁₈	7:3	7.07 ± 0.16	41.08 ± 7.98	1.08 ± 0.08	0.85 ± 0.18	0.57 ± 0.06	1.30 ± 0.09	—	2.13 ± 0.09	37.491	68.766	5.5	0.2943	53.128	1.26	0.0011901	0.309	3.095	6.178 6, 19
(181868) 1999CG ₁₁₉	SDO	7.00 ± 0.04	29.27 ± 11.11	0.87 ± 0.09	0.66 ± 0.12	—	—	—	1.53 ± 0.08	35.182	64.326	16.7	0.2929	49.754	1.74	0.0027276	0.410	2.961	5.770 6
(181871) 1999CO ₅₃	eClas	6.67 ± 0.09	35.09 ± 4.76	1.19 ± 0.15	0.78 ± 0.11	0.56 ± 0.03	1.35 ± 0.10	2.59 ± 0.05	2.00 ± 0.07	39.981	47.767	0.8	0.0887	43.874	0.40	0.0001166	0.090	3.091	5.903 6, 18
(181874) 1999HW ₁₁	SDTNO	6.71 ± 0.06	13.11 ± 4.75	0.83 ± 0.06	0.51 ± 0.08	0.49 ± 0.08	0.99 ± 0.08	1.84 ± 0.04	1.32 ± 0.04	39.069	66.142	17.2	0.2573	52.605	1.62	0.0024523	0.392	3.013	5.969 6, 8, 24
(181902) 1999RD ₂₁₅	SDTNO	—	—	—	—	0.50 ± 0.06	0.88 ± 0.10	—	—	37.663	209.857	25.9	0.6957	123.760	2.22	0.0003958	0.821	2.865	6.345 5
(182222) 2000YU ₁	7:4	6.42 ± 0.06	24.77 ± 3.23	—	0.66 ± 0.06	0.52 ± 0.04	1.18 ± 0.07	—	—	39.577	48.158	6.1	0.0978	43.868	0.65	0.0003939	0.144	3.076	5.865 15
(182223) 2000YC ₂	SDO	7.12 ± 0.11	18.00 ± 6.58	—	0.67 ± 0.11	0.24 ± 0.11	0.91 ± 0.15	—	—	36.133	81.841	19.9	0.3874	58.987	2.01	0.0035454	0.515	2.939	5.927 15
(182397) 2001QW ₂₉₇	9:4	6.66 ± 0.06	24.33 ± 4.03	1.02 ± 0.10	0.58 ± 0.07	0.67 ± 0.06	1.25 ± 0.09	2.27 ± 0.09	1.60 ± 0.07	39.587	63.579	17.1	0.2326	51.583	1.56	0.0023233	0.374	3.019	5.956 24
(182933) 2002GZ ₅₁	SDTNO(b)	—	47.24 ± 4.69	0.94 ± 0.04	0.81 ± 0.05	—	1.35 ± 0.06	—	1.75 ± 0.06	38.312	62.185	1.1	0.2376	50.248	1.00	0.0007086	0.238	3.109	6.140 21
(182934) 2002GL ₁₃₂	hClas	5.47 ± 0.19	29.03 ± 8.89	1.06 ± 0.28	0.65 ± 0.11	0.69 ± 0.30	1.43 ± 0.33	2.42 ± 0.19	1.68 ± 0.26	39.257	49.026	11.6	0.1107	44.142	1.04	0.0010954	0.230	3.040	5.788 22, 15

Notes. ^(a) Dynamical class or family — see Sect. 2. Binary or multiple objects are indicated with (b); retrograde orbits with (r); and Haumea collisional family objects with (h). ^(b) R-filter absolute magnitude not corrected for phase effects. It may also be denoted by $H_R(\alpha)$ or $R(1, \alpha)$. ^(c) Spectral gradient, slope parameter, or reddening S . It expresses the reflectivity spectrum variation in percent of reddening per 1000 Å — see Hainaut & Delsanti (2002). ^(d) Collisional velocity. The mean square of an object's encounter velocity with its circular analog — see Sect. 5.6.2. ^(e) Öpik's ψ . Another estimate of an object's encounter velocity with its circular analog — see Sect. 5.6.2. ^(f) Orbital excitation. It relates with collisional velocity — see Sect. 5.6.2. ^(g) Tisserand parameter, or invariant, relative to Neptune — see Sect. 5.6.2. ^(h) Tisserand parameter, or invariant, relative to Jupiter.

References. (1) Luu & Jewitt (1996); (2) Boehnhardt et al. (2001); (3) Tegler & Romanishin (2000); (4) Gil-Hutton & Licandro (2001); (5) Boehnhardt et al. (2002); (6) Trujillo & Brown (2002); (7) Jewitt & Luu (2001); (8) Benecchi et al. (2011); (9) Green et al. (1997); (10) Barucci et al. (2000); (11) Doressoundiram et al. (2002); (12) Jewitt & Luu (1998); (13) Tegler & Romanishin (2003); (14) Delsanti et al. (2001); (15) Santos-Sanz et al. (2009); (16) Snodgrass et al. (2010); (17) Doressoundiram et al. (2001); (18) Sheppard (2012); (19) Peixinho et al. (2004); (20) Tegler et al. (2003); (21) Gulbis et al. (2006); (22) Doressoundiram et al. (2005b); (23) Doressoundiram et al. (2007); (24) Sheppard (2010); (25) Tegler et al. <http://www.physics.nau.edu/~teglar/research/survey.htm>; (26) Fornasier et al. (2004); (27) Perna et al. (2010); (28) Rabinowitz et al. (2008); (29) Romanishin et al. (2010); (30) Lazzaro et al. (1997); (31) Bauer et al. (2003); (32) Romon-Martin et al. (2003); (33) Romanishin et al. (1997); (34) Romon-Martin et al. (2002); (35) Rabinowitz et al. (2007); (36) Peixinho et al. (2001); (37) Tegler & Romanishin (1997); (38) Barucci et al. (1999); (39) Hainaut et al. (2000); (40) Boehnhardt et al. (2004); (41) Barucci et al. (2002); (42) Ferrin et al. (2001); (43) Alvarez-Candal et al. (2010); (44) Dotto et al. (2003); (45) DeMeo et al. (2010); (46) Doressoundiram et al. (2005a); (47) de Bergh et al. (2005); (48) Rabinowitz et al. (2006); (49) Benecchi et al. (2009); (50) Peixinho et al. (2012)